

# The structure of minimizers of the frame potential on fusion frames <sup>\*†</sup>

Pedro G. Massey, Mariano A. Ruiz and Demetrio Stojanoff<sup>‡</sup>

## Abstract

In this paper we study the fusion frame potential, that is a generalization of the Benedetto-Fickus (vectorial) frame potential to the finite-dimensional fusion frame setting. The structure of local and global minimizers of this potential is studied, when we restrict the frame potential to suitable sets of fusion frames. These minimizers are related to tight fusion frames as in the classical vector frame case. Still, tight fusion frames are not as frequent as tight frames; indeed we show that there are choices of parameters involved in fusion frames for which no tight fusion frame can exist. Thus, we exhibit necessary and sufficient conditions for the existence of tight fusion frames with prescribed parameters, involving the so-called Horn-Klyachko's compatibility inequalities. The second part of the work is devoted to the study of the minimization of the fusion frame potential on a fixed sequence of subspaces, varying the sequence of weights. We related this problem to the index of the Hadamard product by positive matrices and use it to give different characterizations of these minima.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Preliminaries and Notations.</b>	<b>3</b>
2.1	Frames of subspaces, or fusion frames for $\mathbb{C}^n$	4
2.2	Sets of fusion frames and the FF-potential	4
2.3	Klyachko-Fulton approach	6
<b>3</b>	<b>On the existence of tight fusion frames.</b>	<b>7</b>
3.1	Dimensional restrictions	8
3.2	Characterizations for fixed weights	8
<b>4</b>	<b>Minimization for fixed weights</b>	<b>10</b>
4.1	Lower bound for the potential	10
4.2	Structure of local minima: The geometrical approach	12
4.3	The eigenvalues of all $d_S$ -minimizers coincide	16
<b>5</b>	<b>Minimization for fixed subspaces</b>	<b>18</b>
5.1	Minimal weights	18
5.2	Critical Points and local minimizers.	19

---

<sup>\*</sup>Keywords: Fusion frames, frame potential, majorization, Hadamard product.

<sup>†</sup>Mathematics subject classification (2000): 42C15, 15A60.

<sup>‡</sup>Partially supported by CONICET (PIP 5272/05) and Universidad de La PLata (UNLP 11 X472).

<b>A Hadamard products and indexes.</b>	<b>23</b>
A.1 Basic definitions and properties . . . . .	23

# 1 Introduction

Fusion frames were introduced by P. G. Casazza and G. Kutyniok under the name of “frame of subspaces” in [8]. They are a generalization of the usual frames of vectors for a Hilbert space  $\mathcal{H}$ ; indeed frames of vectors can be treated as “one dimensional fusion frames”. During the last years, the theory of fusion frames has been a fast-growing area. Several applications of fusion frames have been studied, for example, sensor networks [11], neurology [19], coding theory [4], [5], [16], among others. Specifically, applications which require distributed processing can be well described and studied using fusion frames. We refer the reader to the work by P.G. Casazza, G. Kutyniok and S. Li [10] and the references therein, for a detailed treatment of the fusion frame theory. Further developments can be found in [7], [9] and [20].

In finite dimension, a fusion frame is a sequence of subspaces of  $\mathbb{F}^n$  ( $\mathbb{F} = \mathbb{C}$  or  $\mathbb{R}$ ) together with a set of positive weights such that the weighted sum of the orthogonal projections to these subspaces (called *fusion frame operator*) is a positive invertible operator (see Definition 2.1.1). As in the case of vector frames, it is usually desired this invertible operator to be a multiple of the identity. In this case, the frame is called a tight fusion frame, also noted TFF. However, tight fusion frames might not exist for a fixed choice of dimensions  $\mathbf{d} = (d_i)_{i \in \mathbb{I}_m}$  of the subspaces, for any sequence of weights  $w = (w_i)_{i \in \mathbb{I}_m}$  (see the discussion following Proposition 3.1.1).

Inspired on the work in classical frame theory (see [2, 18]), we study a convex functional on fusion frame operators (the FF potential or FFP), also studied by Casazza and Fickus in [7], which generalizes the Benedetto-Fickus frame potential. In this paper we analyze its local and global minima. This is motivated by the fact that local (global) minimizers characterize unit norm tight vector frames (see [2, 4, 5, 6, 18]). Since the FF potential can be seen as a “measure of orthogonality” of the frame vectors, it provides also an interesting geometrical description of fusion frames. These considerations motivate the study this type of minimizations in the fusion frame context. A related study can be found in [7], a work that the authors became aware in an advanced stage of writing this paper.

The main tool used in [18] for these problems for vector frames, namely majorization of matrices, can be replaced in the context of fusion frames by the theory developed by Horn and Klyachko in order to have a spectral characterization of hermitian matrices which are the sum of a set of hermitian matrices. For example, this approach provides necessary and sufficient conditions of existence of TFF’s, summarized in a family of inequalities. This technique, although seems to be rather impractical due to the complex conditions involved, becomes an useful tool in the study of the spectral structure of the FF potential minimizers.

We first consider the problem of existence of TFF’s. We show some dimensional restrictions on the subspaces regarding this problem, and then give equivalent conditions for the case of fixed dimensions and weights. We refer to [7] for further developments in this direction.

The rest of the paper deals with the minimization of the FF potential on some sets of Fusion Bessel sequences (i.e. sets of projections and weights whose fusion frame operator is not necessary invertible). Mainly, we work on Bessel sequences with fusion frame operator of trace one. This is a natural restriction in order to avoid scalar multiplications, and it allows an interpretation of the FF potential as a measure of how far is the fusion frame operator to the suitable multiple of the identity corresponding to the (possibly non-existing) tight fusion frames with trace one. A detailed discussion of this approach can be found in subsection 2.2.

The minimization of the FFP is done in three different settings: first, by fixing the weights  $w$  and the dimensions  $\mathbf{d}$  of the subspaces. Then, fixing only the dimensions  $\mathbf{d}$ . Finally, we consider a fixed sequence of subspaces  $(W_i)_{i \in \mathbb{I}_m}$ , and optimize over the set of admissible weights. In the three cases, the minimization is made under the previously mentioned “trace one” restriction.

For the first problem, a geometrical approach similar to that done in [18] allows us to obtain a characterization of local minimizers of the FF potential: they are orthogonal sums of tight frames on each eigenspace of the frame operator. Then, using Horn and Klyachko techniques, we prove that all minimizers (even those which are local minimizers) have the same eigenvalues, with the same multiplicities. Similar results are obtained for the second problem (fixing only the dimensions of the subspaces).

The last section of the paper is devoted to the study of the optimization of the fusion frame potential of fusion frames obtained from a fixed sequence of subspaces in  $\mathbb{C}^n$  which generate the whole space. Since every sequence of weights makes it a fusion frame, we seek for the best choice of weights, meaning those which minimize (globally) the FF potential. Then, we establish a connection between this optimization problem and the *Hadamard indexes* studied in [12] (which involve the Hadamard or entry-wise product of matrices) of a kind of Gramm matrix associated to the fixed subspaces. Using these tools, we get a characterization of the set of optimal weights, and a way to compute them under some reasonable assumptions on the initial sequence of subspaces. This analysis seems to be new even for the case of vector frames.

However, as it is shown by an example, the minimizers could “erase” some of the initial subspaces (i.e. the set of optimal weights could have zeros). Moreover, it is possible to obtain a minimizer which is a Bessel sequence of subspaces which is not generating, a phenomenon which does not happen in the previous settings. Motivated by this problem, we study the geometry of the set of all weights  $w$  which minimize the FF potential, and in particular their possible supports (namely, those sub-indexes  $i$  such that  $w_i > 0$ ). At the end of the section, we present some examples which illustrate these type of anomalies.

The paper is organized as follows: Section 2 contains preliminary definitions on fusion frames and the basic notation used throughout the paper. This section also contains a brief exposition of Horn-Klyachko’s compatibility inequalities. Section 3 is devoted to the study of minimizers of the FF potential, restricted to the sets of fusion frames detailed before. In Section 4 we analyze the problem of minimize the FF potential for a fixed sequence of subspaces, varying the weights. The paper ends with an appendix containing definitions and several results concerning Hadamard indexes of positive matrices, which are related to the contents of Section 4.

We wish to acknowledge to P. Casazza and M. Fickus for letting us know about their excellent work [7], which is closely related with the present paper.

## 2 Preliminaries and Notations.

In this paper  $\mathcal{M}_n(\mathbb{C})$  denotes the algebra of complex  $n \times n$  matrices,  $\mathcal{G}l(n)$  the group of all invertible elements of  $\mathcal{M}_n(\mathbb{C})$ ,  $\mathcal{U}(n)$  the group of unitary matrices,  $\mathcal{M}_n(\mathbb{C})_{sa}$  (resp.  $\mathcal{M}_n(\mathbb{C})_{ah}$ ) denotes the real subspace of hermitian (resp. anti-hermitian) matrices,  $\mathcal{M}_n(\mathbb{C})^+$  the set of positive semidefinite matrices, and  $\mathcal{G}l(n)^+ = \mathcal{M}_n(\mathbb{C})^+ \cap \mathcal{G}l(n)$ . Given  $T \in \mathcal{M}_n(\mathbb{C})$ ,  $R(T)$  denotes the image of  $T$ ,  $N(T)$  the null space of  $T$ ,  $\sigma(T)$  the spectrum of  $T$ ,  $\text{tr } T$  the trace of  $T$ , and  $\text{rk } T$  the rank of  $T$ .

Given  $m \in \mathbb{N}$  we denote by  $\mathbb{I}_m = \{1, \dots, m\}$  and  $\mathbf{1} = \mathbf{1}_m \in \mathbb{R}^m$  denotes the vector with all its entries equal to 1. If  $v \in \mathbb{R}^n$ , we denote by  $\text{diag}(v) \in \mathcal{M}_n(\mathbb{C})$  the diagonal matrix with  $v$  in its diagonal, and by  $v^\downarrow \in \mathbb{R}^n$  the vector obtained by re-arrangement of the coordinates of  $v$  in non-increasing order. If  $T \in \mathcal{M}_n(\mathbb{C})_{sa}$ , we denote by  $\lambda(T) \in \mathbb{R}^n$  the vector of eigenvalues of  $T$ , counted

with multiplicities, in such a way that  $\lambda(T) = \lambda(T)^\perp$ .

Given a subspace  $W \subseteq \mathbb{C}^n$ , we denote by  $P_W \in \mathcal{M}_n(\mathbb{C})^+$  the orthogonal projection onto  $W$ , i.e.  $R(P_W) = W$  and  $N(P_W) = W^\perp$ . For vectors on  $\mathbb{C}^n$  we shall use the euclidean norm, but for matrices  $T \in \mathcal{M}_n(\mathbb{C})$ , we shall use both the spectral norm  $\|T\| = \|T\|_{sp} = \max_{\|x\|=1} \|Tx\|$ , and the

Frobenius norm  $\|T\|_2 = (\text{tr } T^*T)^{\frac{1}{2}} = \left( \sum_{i,j \in \mathbb{I}_n} |T_{ij}|^2 \right)^{\frac{1}{2}}$ . This norm is induced by the inner product  $\langle A, B \rangle = \text{tr } B^*A$ , for  $A, B \in \mathcal{M}_n(\mathbb{C})$ .

## 2.1 Frames of subspaces, or fusion frames for $\mathbb{C}^n$

We begin by defining the basic notions of fusion frame theory in the finite dimensional context. For an introduction to fusion frames for general Hilbert spaces, see [8], [10] or [20]. Briefly, a fusion frame for  $\mathbb{C}^n$  is a generating sequence of subspaces, equipped with weights assigned to each subspace. Nevertheless, we prefer to give the “frame style” definition, which adjusts better to our purposes.

**Definition 2.1.1.** Let  $\mathcal{W} = \{W_i\}_{i \in \mathbb{I}_m}$  be closed subspaces of  $\mathcal{H} \cong \mathbb{C}^n$ , and  $w = \{w_i\}_{i \in \mathbb{I}_m} \in \mathbb{R}_{>0}^m$ . The sequence  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m}$  is a *fusion frame (FF)* for  $\mathcal{H}$ , if there exist  $A, B > 0$  such that

$$A\|f\|^2 \leq \sum_{i \in \mathbb{I}_m} w_i^2 \|P_{W_i} f\|^2 \leq B\|f\|^2 \quad \text{for every } f \in \mathcal{H} . \quad (1)$$

If only the right-hand side inequality in (1) holds, then we say that  $\mathcal{W}_w$  is a *Bessel sequence of subspaces (BSS)* for  $\mathcal{H}$ . The **frame operator** of  $\mathcal{W}_w$  is defined by the formula

$$S_{\mathcal{W}_w} = \sum_{i \in \mathbb{I}_m} w_i^2 P_{W_i} \in \mathcal{M}_n(\mathbb{C})^+ . \quad (2)$$

Observe that  $\mathcal{W}_w$  is a FF if and only if  $S_{\mathcal{W}_w} \in \mathcal{G}l(n)^+$  and, in this case  $A I_n \leq S_{\mathcal{W}_w} \leq B I_n$ . We say that  $\mathcal{W}_w$  is a *tight FF (TFF)* if  $A = B$ , in other words, if  $S_{\mathcal{W}_w} = A I_n$ .  $\triangle$

**Remark 2.1.2.** Let  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m}$  be a BSS for  $\mathbb{C}^n$ . For each  $i \in \mathbb{I}_m$ , we can take an o.n.b.  $\mathcal{B}_i = \{e_j^{(i)}\}_{j \in J_i}$  of  $W_i$ . Hence, for every  $f \in \mathcal{H}$ , we have that

$$P_{W_i} f = \sum_{j \in J_i} \langle f, e_j^{(i)} \rangle e_j^{(i)} , \quad \text{for } i \in \mathbb{I}_m \implies S_{\mathcal{W}_w} f = \sum_{i \in \mathbb{I}_m} w_i^2 P_{W_i} f = \sum_{i \in \mathbb{I}_m} \sum_{j \in J_i} \langle f, w_i e_j^{(i)} \rangle w_i e_j^{(i)} .$$

Therefore,  $\mathcal{W}_w$  induces a vector Bessel sequence  $\mathcal{F} = \{w_i e_j^{(i)} : i \in \mathbb{I}_m, j \in J_i\}$  which has a very useful property: Its frame operator  $S_{\mathcal{F}} = S_{\mathcal{W}_w}$ .  $\triangle$

## 2.2 Sets of fusion frames and the FF-potential

We shall establish several notations regarding sets of FF's and BSS's :

**Notations 2.2.1.** Fix  $n, m \in \mathbb{N}$  and consider a Hilbert space  $\mathcal{H} \cong \mathbb{C}^n$ .

1. We shall denote by  $\mathcal{S}_{m,n}$  the set of all FF's of the form  $\mathcal{W}_w = (w, \mathcal{W}) = (w_i, W_i)_{i \in \mathbb{I}_m}$ , where  $w \in \mathbb{R}_{>0}^m$  and  $\mathcal{W}$  a generating sequence of subspaces of  $\mathcal{H}$ .

2. Given a sequence  $\mathbf{d} \in \mathbb{N}^m$  such that  $\text{tr } \mathbf{d} = \sum_{i \in \mathbb{I}_m} d_i \geq n$ , we denote by

$$\mathcal{S}_{m,n}(\mathbf{d}) = \{ \mathcal{W}_w \in \mathcal{S}_{m,n} : \dim W_i = d_i \text{ for every } i \in \mathbb{I}_m \}, \quad (3)$$

Similarly, we denote by  $\mathcal{B}_{m,n}(\mathbf{d})$  the set of BSS's with the same dimensional restrictions.

3. Given  $v \in \mathbb{R}_{>0}^m$ , we denote by

$$\mathcal{B}_{m,n}(\mathbf{d}, v) = \{ \mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}) : w = v \} \quad \text{and} \quad \mathcal{S}_{m,n}(\mathbf{d}, v) = \mathcal{B}_{m,n}(\mathbf{d}, v) \cap \mathcal{S}_{m,n}(\mathbf{d}), \quad (4)$$

the subsets of  $\mathcal{B}_{m,n}(\mathbf{d})$  and  $\mathcal{S}_{m,n}(\mathbf{d})$  with a fixed sequence of weights  $v$ .

4. Finally, we denote by

$$\mathcal{B}_{m,n}^1(\mathbf{d}) = \{ \mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}) : \text{tr } S_{\mathcal{W}_w} = \sum_{i \in \mathbb{I}_m} w_i^2 d_i = 1 \} \quad . \quad (5)$$

and  $\mathcal{S}_{m,n}^1(\mathbf{d}) = \mathcal{B}_{m,n}^1(\mathbf{d}) \cap \mathcal{S}_{m,n}(\mathbf{d})$ .

5. We say that a pair  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  is **normalized** if  $\text{tr } \mathbf{d} \geq n$  and  $\sum_{i \in \mathbb{I}_m} w_i^2 d_i = 1$ . Observe that  $(\mathbf{d}, w)$  is normalized if and only if  $\mathcal{B}_{m,n}(\mathbf{d}, w) \subseteq \mathcal{B}_{m,n}^1(\mathbf{d})$ .  $\triangle$

The following definition is suggested by the classical Benedetto-Fickus potential, whose value in a vector frame  $\mathcal{F}$  can be calculated as  $\text{FP}(\mathcal{F}) = \text{tr } S_{\mathcal{F}}^2$ .

**Definition 2.2.2.** Given a BSS  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m}$ , the Benedetto-Fickus *fusion frame potential* (FF-potential) of  $\mathcal{W}_w$  is given by:

$$\text{FFP}(\mathcal{W}_w) = \sum_{i,j=1}^m w_i^2 w_j^2 \text{tr}(P_{W_i} P_{W_j}) = \text{tr } S_{\mathcal{W}_w}^2 \quad . \quad (6)$$

Notice that in this case  $\text{FFP}(\mathcal{W}_w) = \text{FP}(\mathcal{F})$ , for any vector Bessel sequence  $\mathcal{F}$  obtained from  $\mathcal{W}_w$  as in Remark 2.1.2. We define also the following matrix:

$$\mathbf{P}_q(\mathcal{W}_w) = \sum_{i,j=1}^m w_i^2 w_j^2 |P_{W_i} P_{W_j}|^2 = \sum_{i,j=1}^m w_i^2 w_j^2 P_{W_j} P_{W_i} P_{W_j} \in \mathcal{M}_n(\mathbb{C})^+ \quad . \quad (7)$$

The matrix  $\mathbf{P}_q(\cdot)$  is related to the so-called q-potential [17] defined in the more general context of reconstruction systems. Notice that the Benedetto-Fickus fusion frame potential can be computed in terms of  $\mathbf{P}_q(\mathcal{W}_w)$ , since  $\text{FFP}(\mathcal{W}_w) = \text{tr } \mathbf{P}_q(\mathcal{W}_w)$ .  $\triangle$

The scope of this paper is to study minimizers of the FF-potential. In order to avoid scalar multiplications (note that  $\text{FFP}(\mathcal{W}_{t \cdot w}) = t^4 \text{FFP}(\mathcal{W}_w)$ ) we shall restrict ourselves to minimize the FF-potential on subsets of  $\mathcal{S}_{m,n}^1(\mathbf{d})$  or  $\mathcal{B}_{m,n}^1(\mathbf{d})$ , for  $\mathbf{d}$  and  $m$  fixed. In other words, we shall minimize the FF-potential for those frames  $\mathcal{W}_w$  such that  $\text{tr } S_{\mathcal{W}_w} = 1$ .

This specific restriction is justified because, if there exist tight FF's in  $\mathcal{S}_{m,n}(\mathbf{d})$ , then their FF-potential and frame bounds are determined exactly by the trace of their frame operators. Namely, if  $\mathcal{W}_w \in \mathcal{S}_{m,n}(\mathbf{d})$  is tight, and  $\text{tr } S_{\mathcal{W}_w} = a$ , then  $S_{\mathcal{W}_w} = \frac{a}{n} I_n$  and  $\text{FFP}(\mathcal{W}_w) = \frac{a^2}{n}$ .

Even in the case that there are no TFF's in  $\mathcal{S}_{m,n}(\mathbf{d})$ , this restriction seems to be quite natural. Indeed, for BSS's with fixed trace, the FF-potential can be seen as a measure of the (Frobenius) distance of their frame operators to a **fixed** multiple of the identity:

**Proposition 2.2.3.** *Let  $\mathcal{W}_w \in \mathcal{S}_{m,n}^1(\mathbf{d})$ . Then*

$$\left\| \frac{1}{n} I_n - S_{\mathcal{W}_w} \right\|_2^2 = \text{tr } S_{\mathcal{W}_w}^2 - \frac{1}{n} = \text{FFP}(\mathcal{W}_w) - \frac{1}{n} .$$

*Proof.* Since  $\text{tr } S_{\mathcal{W}_w} = 1$ , a direct computation shows that

$$\left\| \frac{1}{n} I_n - S_{\mathcal{W}_w} \right\|_2^2 = \text{tr} \left( \frac{1}{n^2} I_n - \frac{2}{n} S_{\mathcal{W}_w} + S_{\mathcal{W}_w}^2 \right) = \text{tr } S_{\mathcal{W}_w}^2 - \frac{1}{n} . \quad \square$$

The last result shows that if there exist tight FF's in  $\mathcal{S}_{m,n}^1(\mathbf{d})$ , then they are the unique global minimizers of the FF-potential on  $\mathcal{S}_{m,n}^1(\mathbf{d})$ . But in the case that there are no TFF's in  $\mathcal{S}_{m,n}^1(\mathbf{d})$ , the minimization of the FF-potential becomes more interesting: it provides the elements of  $\mathcal{S}_{m,n}^1(\mathbf{d})$  that can be expected to have the best properties.

In this paper we deal mostly with these type of minimizations under two different further restrictions: we work in the set  $\mathcal{B}_{m,n}(\mathbf{d}, w) \subseteq \mathcal{B}_{m,n}^1(\mathbf{d})$  for a fixed normalized pair  $(\mathbf{d}, w)$ , or we fix a generating sequence  $\mathcal{W}$  of subspaces, and minimize the FF-potential over all sequences  $w \in \mathbb{R}_{\geq 0}^m$  such that  $\mathcal{W}_w \in \mathcal{B}_{m,n}^1(\mathbf{d})$ .

## 2.3 Klyachko-Fulton approach

Recall that given  $x \in \mathbb{R}^n$ , we denote by  $x^\downarrow \in \mathbb{R}^n$  the vector obtained by re-arrangement of the coordinates of  $x$  in non-increasing order. Given  $x, y \in \mathbb{R}^n$  we say that  $x$  is *submajorized* by  $y$ , and write  $x \prec_w y$ , if  $\sum_{i=1}^k x_i^\downarrow \leq \sum_{i=1}^k y_i^\downarrow$  for every  $k \in \mathbb{I}_n$ . If we further have that  $\text{tr}(x) := \sum_{i=1}^n x_i = \sum_{i=1}^n y_i$  then we say that  $x$  is *majorized* by  $y$ , and write  $x \prec y$ .

**Example 2.3.1.** As an elementary example, that we shall use in what follows, let  $x \in \mathbb{R}_{\geq 0}^n$  and  $0 \leq a \leq \text{tr}(x) \leq b$ . The reader can easily verify that  $\frac{a}{n} \mathbf{1}_n \prec_w x \prec_w b e_1$ .  $\triangle$

(Sub)majorization between vectors is extended by T. Ando in [1] to (sub)majorization between self-adjoint matrices as follows : given  $A, B \in \mathcal{M}_n(\mathbb{C})_{sa}$ , we say that  $A$  is *submajorized* by  $B$ , and write  $A \prec_w B$ , if  $\lambda(A) \prec_w \lambda(B)$ . If we further have that  $\text{tr}(A) = \text{tr}(B)$  then we say that  $A$  is *majorized* by  $B$  and write  $A \prec B$ .

Although simple, submajorization plays a central role in optimization problems with respect to convex functionals and unitarily invariant norms, as the following result shows (for a detailed account in majorization see Bhatia's book [3]).

**Theorem 2.3.2.** Let  $A, B \in \mathcal{M}_n(\mathbb{C})^{sa}$ . Then, the following statements are equivalent:

1.  $A \prec_w B$ .
2. For every unitarily invariant norm  $\|\cdot\|$  in  $\mathcal{M}_n(\mathbb{C})$  we have  $\|A\| \leq \|B\|$ .
3. For every increasing convex function  $f : \mathbb{R} \rightarrow \mathbb{R}$  we have  $\text{tr } f(A) \leq \text{tr } f(B)$ .

Moreover, if  $A \prec_w B$  and there exists an increasing strictly convex function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $\text{tr } f(A) = \text{tr } f(B)$  then there exists  $U \in \mathcal{U}(n)$  such that  $A = U^* B U$ .  $\square$

In what follows we describe the basic facts about the spectral characterization of the sums of hermitian matrices obtained by Klyachko [15] and Fulton [14]. Let

$$\mathcal{K}_r^n = \{(j_1, \dots, j_r) \in \mathbb{I}_n^r : j_1 < j_2 < \dots < j_r\}.$$

For  $J = (j_1, \dots, j_r) \in \mathcal{K}_r^n$ , define the associated partition

$$\lambda(J) = (j_r - r, \dots, j_1 - 1).$$

Denote by  $LR_r^n(m)$  the set of  $(m+1)$ -tuples  $(J_0, \dots, J_m) \in (\mathcal{K}_r^n)^{m+1}$ , such that the Littlewood-Richardson coefficient of the associated partitions  $\lambda(J_0), \dots, \lambda(J_m)$  is positive, i.e. one can generate the Young diagram of  $\lambda(J_0)$  from those of  $\lambda(J_1), \dots, \lambda(J_m)$  according to the Littlewood-Richardson rule (see [13]). With these notations and terminologies we have

**Theorem 2.3.3** ([15, 14]). Let  $\lambda_i = \lambda_i^\downarrow = (\lambda_1^{(i)}, \dots, \lambda_n^{(i)}) \in \mathbb{R}^n$  for  $i = 0, \dots, m$ . Then, the following statements are equivalent:

1. There exists  $A_i \in \mathcal{M}_n(\mathbb{C})_{sa}$  with  $\lambda(A_i) = \lambda_i$  for  $0 \leq i \leq m$  and such that

$$A_0 = A_1 + \dots + A_m.$$

2. For each  $r \in \{1, \dots, n\}$  and  $(J_0, \dots, J_m) \in LR_r^n(m)$  we have

$$\sum_{j \in J_0} \lambda_j^{(0)} \leq \sum_{i=1}^m \sum_{j \in J_i} \lambda_j^{(i)} \quad (8)$$

plus the condition  $\sum_{j=1}^n \lambda_j^{(0)} = \sum_{i=1}^m \sum_{j=1}^n \lambda_j^{(i)}.$

Moreover, if  $(A_i)_{i=0}^m$  are as in item 1. above and  $(J_0, \dots, J_m) \in LR_r^n(m)$  satisfy equality in (8), then there exists a subspace  $L \subseteq \mathbb{C}^n$  with  $\dim L = r$ , that simultaneously reduces  $A_i$  for  $0 \leq i \leq m$  and such that  $\lambda(P_L A_i) = (\lambda_j^{(i)})_{j \in J_i}$ , where  $P_L$  denotes the orthogonal projection of  $\mathbb{C}^n$  onto  $L$ .  $\square$

We shall refer to the inequalities in (8) as *Horn-Klyachko's compatibility inequalities*.

### 3 On the existence of tight fusion frames.

The following facts exemplify the difference between the theory of vector frames and that of frames of subspaces. In [2] (see also [6] and [18]) it is shown that the local minimizers of the frame potential on the set

$$F_{m,n}^1 = \{\mathcal{F} = \{f_i\}_{i \in \mathbb{I}_m} : \text{each } f_i \in \mathbb{C}^n \text{ and } \text{tr}(S_{\mathcal{F}}) = \sum_{i \in \mathbb{I}_m} \|f_i\|^2 = 1\}$$

are tight frames. Since the set  $F_{m,n}^1$  is compact and the frame potential is a continuous function, there must be global (and hence local) minima of the frame potential. This was used to give an indirect proof of the existence of such frames in the vectorial case.

### 3.1 Dimensional restrictions

Let  $\mathbf{d} \in \mathbb{N}^m$  with  $\text{tr } \mathbf{d} \geq n$  and consider the set  $\mathcal{S}_{m,n}^1(\mathbf{d})$  defined in Eq. (5). Using Remark 2.1.2 it follows that if  $\mathbf{d} = \mathbf{1}_m$ , then we can identify  $\mathcal{S}_{m,n}^1(\mathbf{d})$  with  $F_{m,n}^1$ , and the previous comments can be applied. Hence it seems natural to ask whether there always exist TFF's in  $\mathcal{S}_{m,n}^1(\mathbf{d})$ , since they would be all the global minimizers of the FF-potential on  $\mathcal{S}_{m,n}^1(\mathbf{d})$ . The following results show that in general the answer is no.

**Proposition 3.1.1.** *Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair, with  $M = \text{tr } \mathbf{d} \geq n$ , and assume that  $\mathcal{W}_w \in \mathcal{S}_{m,n}^1(\mathbf{d})$  is a TFF, so that  $S_{\mathcal{W}_w} = \frac{1}{n} I_n$ . If there exists  $i \in \mathbb{I}_m$  such that*

$$M - d_i = \sum_{k \neq i} d_k \leq n - 1 \implies w_i^2 = \frac{1}{n} \quad \text{and} \quad P_{W_i} P_{W_j} = 0 \quad \text{for every } j \in \mathbb{I}_m \setminus \{i\}.$$

*Proof.* Consider the tight vector frame  $\mathcal{F} = \{w_i^2 e_j^{(i)} : i \in \mathbb{I}_m, j \in \mathbb{I}_{d_i}\}$  associated to  $\mathcal{W}_w$ , as described in Remark 2.1.2. Let  $G \in \mathcal{M}_M(\mathbb{C})^+$  denote the Gramm matrix of the vector frame  $\mathcal{F}$  and let  $G_i = w_i^2 I_{d_i}$  denote the Gramm matrix of each subsequence  $\{w_i^2 e_j^{(i)} : j \in \mathbb{I}_{d_i}\}$ . Then each  $G_i$  is a  $d_i \times d_i$  principal sub-matrix of  $G$ . By Cauchy's interlacing principle [3] we get:

$$\lambda_j(G) \geq \lambda_j(G_i) \geq \lambda_{M-d_i+j}(G) \quad \text{for } 1 \leq j \leq d_i,$$

where  $\lambda(G) = (\lambda_j(G))_{j \in \mathbb{I}_M}$  (resp  $\lambda(G_i) \in \mathbb{R}^{d_i}$ ) denotes the vector of eigenvalues of  $G$  (resp.  $G_i$ ) counting multiplicities and with its entries arranged in non-increasing order. By assumption,

$$\lambda_j(G) = \frac{1}{n} \quad \text{for } j \in \mathbb{I}_n, \quad \lambda_j(G) = 0 \quad \text{for } n < j \leq M, \quad \text{and} \quad \lambda_j(G_i) = w_i^2 \quad \text{for } j \in \mathbb{I}_{d_i}.$$

Thus if  $\sum_{k \neq i} d_k \leq n - 1$ , then  $M - d_i + 1 \leq n$  and  $\frac{1}{n} = \lambda_{M-d_i+1}(G) \leq \lambda_1(G_i) = w_i^2 \leq \lambda_1(G) = \frac{1}{n}$ . It

is known that, in this case, each of the vectors  $e_j^{(i)}$ ,  $1 \leq j \leq d_i$ , must be orthogonal to every other vector in the  $\frac{1}{n}$ -tight vector frame  $\mathcal{F}$ , which implies the last assertion of the theorem.  $\square$

**Example 3.1.2** (About the existence of tight frames in  $\mathcal{S}_{m,n}^1(\mathbf{d})$ ). Consider now  $\mathbf{d} = (2, 2)$  and assume that there exists  $\mathcal{W}_w \in \mathcal{S}_{2,3}^1(\mathbf{d})$  that is tight. That is, we assume that there exist two subspaces  $W_i \subset \mathbb{C}^3$  with  $\dim W_i = 2$ ,  $i = 1, 2$  and  $w_1, w_2 \in \mathbb{R}_{>0}$  such that  $\frac{1}{3} I_3 = w_1^2 P_{W_1} + w_2^2 P_{W_2}$ . Since  $d_1, d_2 \leq 3 - 1$  we conclude from Proposition 3.1.1 that  $w_1^2 = w_2^2 = \frac{1}{3}$ , and  $P_{W_1} P_{W_2} = 0$ , which is impossible. This argument can be extended to show that if the choices of  $\mathbf{d} = (d_i)_{i \in \mathbb{I}_m}$  are such that each  $d_i$  is relatively small compared with  $n$  and  $\sum_{k \neq i} d_k$  then there are no TFF's in

$\mathcal{S}_{m,n}^1(\mathbf{d})$ . For example, in  $\mathcal{S}_{k,2k-1}^1(2 \cdot \mathbf{1}_k)$ ,  $\mathcal{S}_{3,7}^1(3, 3, 3)$ ,  $\mathcal{S}_{3,9}^1(4, 4, 4)$ , etc, there are no TFF's.  $\triangle$

**Remark 3.1.3.** The previous results show some dimensional restrictions for the existence of TFF's in  $\mathcal{S}_{m,n}^1(\mathbf{d})$ . In the paper by Casazza and Fickus [7], some sufficient conditions on  $n$ ,  $m$  and  $\mathbf{d}$  are given (particularly if  $\mathbf{d}$  is a multiple of  $\mathbf{1}_m$ ), which assure the existence in  $\mathcal{S}_{m,n}^1(\mathbf{d})$  of such fusion frames. For further results in this direction, see also [16].  $\triangle$

### 3.2 Characterizations for fixed weights

The following theorem gives us some general bounds for  $\mathbf{P}_q(\cdot)$  and states several conditions on  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d})$  which are equivalent to the assertion that  $\mathcal{W}_w$  is a  $\frac{1}{n}$ -TFF.



**Theorem 3.2.1.** Let  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d})$ , where  $\sum_{i \in \mathbb{I}_m} w_i^2 d_i \geq 1$ . Then

$$\frac{1}{n^2} I \prec_w \mathbf{P}_q(\mathcal{W}_w) . \quad (9)$$

For every u.i.n.  $\|\cdot\|$  on  $\mathcal{M}_n(\mathbb{C})$  with associated symmetric gauge function  $\psi$  we have that

$$\frac{1}{n^2} \psi(\mathbf{1}) \leq \|\mathbf{P}_q(\mathcal{W}_w)\| . \quad (10)$$

For every increasing convex function  $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$  with  $f(0) = 0$  we have

$$n \cdot f\left(\frac{1}{n^2}\right) \leq \text{tr } f(\mathbf{P}_q(\mathcal{W}_w)) . \quad (11)$$

Finally, the following conditions are equivalent:

1.  $\mathcal{W}_w$  is a  $\frac{1}{n}$ -TFF.
2. Majorization holds in (9).
3. There exists u.i.n.  $\|\cdot\|$  such that equality holds in (10)
4. There exists an increasing strictly convex function  $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  with  $f(0) = 0$  such that equality holds in (11).

*Proof.* Since  $\text{tr}(S_{\mathcal{W}_w}) = \sum_{i \in \mathbb{I}_m} w_i^2 d_i \geq 1$  then (see Example 2.3.1) it follows that  $\frac{1}{n} I_n \prec_w S_{\mathcal{W}_w}$  and hence

$$\frac{1}{n} = \text{tr}\left(\frac{1}{n} I_n\right)^2 \leq \text{tr } S_{\mathcal{W}_w}^2 = \text{tr } \mathbf{P}_q(\mathcal{W}_w) \implies \frac{1}{n^2} I_n \prec_w \mathbf{P}_q(\mathcal{W}_w) . \quad (12)$$

Notice that by Theorem 2.3.2 then (10) and (11) are consequences of this last fact. Assume that majorization holds in (9), so then we have

$$\text{tr } \frac{1}{n^2} I_n = \text{tr } \mathbf{P}_q(\mathcal{W}_w) = \text{tr } S_{\mathcal{W}_w}^2 .$$

Since  $\frac{1}{n} I_n \prec_w S_{\mathcal{W}_w}$  and the function  $f(x) = x^2$  is strictly convex, by Theorem 2.3.2 we conclude that there exists a unitary  $U \in \mathcal{U}(n)$  such that  $S_{\mathcal{W}_w} = U^* \left(\frac{1}{n} I_n\right) U = \frac{1}{n} I_n$ . On the other hand, if there exists an u.i.n.  $\|\cdot\|$  such that equality holds in (10) then, using the right-hand side of (12) we get

$$\frac{1}{n^2} \psi(\mathbf{1}) = \|\mathbf{P}_q(\mathcal{W}_w)\| \geq \frac{\text{tr } \mathbf{P}_q(\mathcal{W}_w)}{n} \psi(\mathbf{1}) \geq \frac{1}{n^2} \psi(\mathbf{1}) , \quad (13)$$

which implies that  $\text{tr} \left(\frac{1}{n} I_n\right)^2 = \text{tr } \mathbf{P}_q(\mathcal{W}_w)$ . As before, we conclude that  $S_{\mathcal{W}_w} = \frac{1}{n} I_n$ . Similarly, if there exists an increasing strictly convex function  $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  with  $f(0) = 0$  such that equality holds in (11) then Theorem 2.3.2 and the right-hand side of (12) imply that  $S_{\mathcal{W}_w} = \frac{1}{n} I_n$ . Finally, it is clear that in case  $\mathcal{W}_w$  is a TFF then  $\mathbf{P}_q(\mathcal{W}_w) = \frac{1}{n} I_n$ . The last part of the theorem follows from this fact.  $\square$

**Theorem 3.2.2.** Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. Then, the following statements are equivalent:

1. There exists  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m} \in \mathcal{S}_{m,n}^1(\mathbf{d})$  which is a  $\frac{1}{n}$ -TFF.

2. For every  $1 \leq r \leq n-1$  and every  $(J_0, \dots, J_m) \in LR_r^n(m)$  we have that

$$\frac{r}{n} \leq \sum_{i \in \mathbb{I}_m} w_i^2 \cdot |J_i \cap \{1, \dots, d_i\}|.$$

3. There exists an orthogonal projection  $P \in \mathcal{M}_m(\mathcal{M}_n(\mathbb{C}))$  with  $\text{tr}(P) = n$  and such that, if  $\frac{1}{n}P = (w_i w_j P_{ij})_{i,j \in \mathbb{I}_m}$  with  $P_{ij} \in \mathcal{M}_n(\mathbb{C})$  for  $i, j \in \mathbb{I}_m$ , then

$$\frac{1}{w_i^2 n} P_{ii} = \left( \frac{1}{w_i^2 n} P_{ii} \right)^* = \left( \frac{1}{w_i^2 n} P_{ii} \right)^2 \quad \text{and} \quad \text{tr} \left( \frac{1}{w_i^2 n} P_{ii} \right) = d_i \quad \text{for every } i \in \mathbb{I}_m.$$

*Proof.* Notice that the condition in (1) is equivalent to the existence of orthogonal projections  $\{P_i\}_{i \in \mathbb{I}_m}$  such that  $\text{tr}(P_i) = d_i$  for  $1 \leq i \leq m$  and such that  $\sum_{i \in \mathbb{I}_m} w_i^2 P_i = \frac{1}{n} I$ . Hence, by Theorem 2.3.3 it follows that condition (2) should hold, since these are Horn-Klyachko's compatibility inequalities for the spectra of  $\{w_i^2 P_i\}_{i \in \mathbb{I}_m}$  and  $\frac{1}{n} I_n$ . The converse of the previous implication also follows from Theorem 2.3.3 since a self adjoint  $A$  operator with  $\lambda(A) = (\alpha, \dots, \alpha, 0, \dots, 0) \in \mathbb{R}^n$  is necessarily of the form  $A = \alpha P$  for some projection  $P \in \mathcal{M}_n(\mathbb{C})$ .

Assume now (1) and let  $P_i = P_{W_i}$  for  $1 \leq i \leq m$ , where  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m}$  is a  $\frac{1}{n}$ -TFF. Let us consider  $V_i \in \mathcal{M}_n(\mathbb{C})$  a partial isometry such that  $V_i^* V_i = P_i$ ,  $1 \leq i \leq m$ . Define  $V^* = [w_1 V_1^* | w_2 V_2^* | \dots | w_m V_m^*] \in \mathcal{M}_{n, m \cdot n}(\mathbb{C})$  and notice that  $V^* V = \sum_{i \in \mathbb{I}_m} w_i^2 V_i^* V_i = \frac{1}{n} I_n$ . Hence  $V V^* = (w_i w_j V_i V_j^*)_{i,j=1}^m \in \mathcal{M}_m(\mathcal{M}_n(\mathbb{C}))$  is such that  $V V^* = \frac{1}{n} P$  for an orthogonal projection  $P = (P_{ij})_{i,j} \in \mathcal{M}_m(\mathcal{M}_n(\mathbb{C}))$ . By comparing the diagonal blocks we get that

$$\frac{1}{n} P_{ii} = w_i^2 V_i V_i^* \implies \frac{1}{w_i^2 n} P_{ii} = V_i V_i^*, \quad \text{for every } i \in \mathbb{I}_m.$$

Conversely, if  $P = (P_{ij}) \in \mathcal{M}_m(\mathcal{M}_n(\mathbb{C}))$  is as in (3) then there exist matrices  $V_i \in \mathcal{M}_n(\mathbb{C})$  such that if  $V^* = [w_1 V_1^* | \dots | w_m V_m^*] \in \mathcal{M}_{n, m \cdot n}(\mathbb{C})$  then  $P = V V^*$  (since  $\text{rank } P = n$ ). But then, comparing the block diagonal entries we get that  $\frac{1}{n} V_i^* V_i$  is an orthogonal projection with  $\text{tr}(\frac{1}{n} V_i^* V_i) = d_i$  for  $1 \leq i \leq m$ . Hence  $V^* V = I_n$ , that is

$$\sum_{i \in \mathbb{I}_m} w_i^2 \frac{1}{n} V_i^* V_i = \frac{1}{n} I_n.$$

Therefore, if we let  $W_i$  be the range of  $V_i^*$  we get that  $P_{W_i} = \frac{1}{n} V_i^* V_i$  so then  $\mathcal{W} = (w_i, W_i)_{i \in \mathbb{I}_m}$  is a  $\frac{1}{n}$ -TFF for  $\mathbb{C}^n$ .  $\square$

## 4 Minimization for fixed weights

### 4.1 Lower bound for the potential

In this subsection we translate, using Remark 2.1.2, some well known results about vector frames (see [6] or [20]) to the FF context. An interesting fact is that there is a notion of irregularity, defined in terms of the parameters of a given FF, which agree with the vectorial  $n$ -irregularity of their associated vector frames. Nevertheless, the lower bound obtained for the FF-potential is not always attained in the set  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  (see Example 4.1.3).

**Definition 4.1.1.** Given a pair  $\mathbf{d} \in \mathbb{N}^m$  and  $w = w^\perp \in \mathbb{R}_{>0}^m$ , consider its  $q$ -irregularity defined as

$$J_0(\mathbf{d}, w) = \max \left\{ j \in \mathbb{I}_m : \left( n - \sum_{i=1}^j d_i \right) w_j^2 > \sum_{i=j+1}^m w_i^2 d_i \right\},$$

if this set is not empty, otherwise  $J_0(\mathbf{d}, w) = 0$ .

**Proposition 4.1.2.** Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair, where  $w = w^\perp$ . Recall that

$$M = \sum_{i \in \mathbb{I}_m} d_i \geq n. \text{ Let } j_0 := J_0(\mathbf{d}, w) \text{ and } c = \frac{\sum_{i=j_0+1}^m w_i^2 d_i}{n - \sum_{i=1}^{j_0} d_i} < w_{j_0}^2. \text{ If } \mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w) \text{ then}$$

$$FFP(\mathcal{W}_w) \geq \sum_{i=1}^{j_0} d_i w_i^4 + \left( n - \sum_{i=j_0+1}^m d_i \right) c^2. \quad (14)$$

Moreover, equality holds in (14) if and only if the following two conditions hold:

1.  $P_{W_i} P_{W_j} = 0$  for  $1 \leq i \neq j \leq j_0$  and
2.  $\{w_i, W_i\}_{i=j_0+1}^m$  is a TFF for  $\text{span}\{W_i : 1 \leq i \leq j_0\}^\perp$ .

*Proof.* 1. Let  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  and let  $\mathcal{F} = \{w_i^2 e_j^{(i)} : i \in \mathbb{I}_m, j \in \mathbb{I}_{d_i}\}$  be an associated vector frame, as described in Remark 2.1.2. Let  $\mathbf{a} \in \mathbb{R}^M$  denote the vector whose coordinates are the norms of the elements of  $\mathcal{F}$  arranged in non-increasing order. Then,

$$w_k = a_{M(k,j)}, \quad \text{where } M(k,j) = \sum_{i=1}^{k-1} d_i + j, \quad \text{for } k \in \mathbb{I}_m \text{ and } j \in \mathbb{I}_{d_k}.$$

We now consider the  $n$ -irregularity  $r_n(\mathbf{a})$  of the vector  $\mathbf{a}$ :

$$r_n(\mathbf{a}) = \max \left\{ j \in \mathbb{I}_{n-1} : (n-j)a_j > \sum_{i=j+1}^M a_i \right\},$$

if the set on the right is non empty, and  $r_n(\mathbf{a}) = 0$  otherwise. It is straightforward that  $r_n(\mathbf{a}) = \sum_{i=1}^{j_0} d_i$  in the first case. Therefore, inequality (14) can be deduced from [6, Theorem 10] (see also [20]). The same result of [6] shows that equality in Eq. (14) implies that  $S_1 = \{e_j^{(i)} : 1 \leq i \leq j_0, j \in \mathbb{I}_{d_i}\}$  is an orthonormal system in  $\mathbb{C}^n$ , and  $S_2 = \{e_j^{(i)} : j_0 + 1 \leq i \leq m, j \in \mathbb{I}_{d_i}\}$  is a tight frame for  $\mathcal{S}^\perp$ , where  $\mathcal{S} = \text{span}\{S_1\} = \text{span}\{W_i : 1 \leq i \leq j_0\}$ .  $\square$

The following example shows that the lower bound in (14) is not sharp in general.

**Example 4.1.3.** If we set  $n = 3$ ,  $\mathbf{d} = (2, 2)$  and  $w_1 = w_2 = \frac{1}{2}$  then  $\sum_{i \in \mathbb{I}_m} w_i^2 d_i = 1$  and  $J_0(\mathbf{d}, w) = 0$ .

Therefore, by Theorem 4.1.2 the equality (14) holds only in tight fusion frames. Still, there are no tight FS in  $S_{2,3}(\mathbf{d}, w) \subset S_{2,3}(\mathbf{d})$  since the Example 3.1.2 shows that there are no tight FS in the (bigger set)  $S_{2,3}(\mathbf{d})$ .  $\triangle$

## 4.2 Structure of local minima: The geometrical approach

In what follows we consider a perturbation result for Bessel sequences of subspaces. We begin by considering some well known facts from differential geometry that we shall need below. In what follows we consider the unitary group  $\mathcal{U}(n)$  together with its natural differential geometric (Lie) structure. It is well known that the tangent space  $\mathcal{T}_{I_n} \mathcal{U}(n)$  at the identity can be naturally identified with the real vector space

$$\mathcal{M}_n(\mathbb{C})_{ah} = i \cdot \mathcal{M}_n(\mathbb{C})_{sa} = \{X \in \mathcal{M}_n(\mathbb{C}) : X^* = -X\} ,$$

of anti-hermitian matrices. Given  $G \in \mathcal{M}_n(\mathbb{C})^+$  we consider the smooth map

$$\Psi_G : \mathcal{U}(n) \rightarrow \mathcal{U}(G) \subseteq \mathcal{M}_n(\mathbb{C}) \quad \text{given by} \quad \Psi_G(U) = U^* G U , \quad U \in \mathcal{U}(n) , \quad (15)$$

where  $\mathcal{U}(G)$  is the unitary orbit of  $G$ . Under the previous identification, the differential of  $\Psi_G$  at a the point  $I_n \in \mathcal{U}(n)$  in the direction given by  $X \in \mathcal{M}_n(\mathbb{C})_{ah}$  is given by

$$(D\Psi_G)_{I_n}(X) = XG - GX = [X, G] . \quad (16)$$

It is well known that the map  $\Psi_G$  is a submersion of  $\mathcal{U}(n)$  onto  $\mathcal{U}(G)$ . Therefore, the differential  $(D\Psi_G)_{I_n}$  is an epimorphism, and hence (16) gives us a description of the tangent space of the manifold  $\mathcal{U}(G)$  at the point  $G$ : We have that  $T_G \mathcal{U}(G) = \{[X, G] : X \in \mathcal{M}_n(\mathbb{C})_{ah}\}$ .

Let us fix some notations. We denote by

$$\mathcal{M}_n^1(\mathbb{C})_{sa} = \{A \in \mathcal{M}_n(\mathbb{C})_{sa} : \text{tr } A = 1\} \quad \text{and} \quad \mathcal{M}_n^0(\mathbb{C})_{sa} = \{A \in \mathcal{M}_n(\mathbb{C})_{sa} : \text{tr } A = 0\} .$$

Observe that  $\mathcal{M}_n^1(\mathbb{C})_{sa}$  is an affine manifold contained in the real vector space  $\mathcal{M}_n(\mathbb{C})_{sa}$ , whose tangent space is the subspace  $\mathcal{M}_n^0(\mathbb{C})_{sa}$ . On the other hand, given  $X, Y \in \mathcal{M}_n(\mathbb{C})_{sa}$ , it is easy to see that  $\text{tr } XY \in \mathbb{R}$ . Therefore the inner product  $\langle A, B \rangle = \text{tr } B^* A$  of  $\mathcal{M}_n(\mathbb{C})$  still works as a real inner product on  $\mathcal{M}_n(\mathbb{C})_{sa}$ .

Given a set  $\{P_j : j \in \mathbb{I}_m\} \subseteq \mathcal{M}_n(\mathbb{C})_{sa}$  of projections, we denote by

$$\{P_j : j \in \mathbb{I}_m\}' = \{A \in \mathcal{M}_n(\mathbb{C}) : AP_j = P_j A \quad \text{for every} \quad j \in \mathbb{I}_m\} . \quad (17)$$

Note that  $\{P_j : j \in \mathbb{I}_m\}'$  is a closed selfadjoint subalgebra of  $\mathcal{M}_n(\mathbb{C})$ . Therefore, the algebra  $\{P_j : j \in \mathbb{I}_m\}' \neq \mathbb{C} I_n \iff$  there exists a non-trivial orthogonal projection  $Q \in \{P_j : j \in \mathbb{I}_m\}'$ .

**Theorem 4.2.1.** *Let  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m} \in \mathcal{B}_{m,n}^1(\mathbf{d}, w)$  be a BSS. Denote  $P_j = P_{W_j}$  for every  $j \in \mathbb{I}_m$ . Let  $\Psi : \mathcal{U}(n)^m \rightarrow \mathcal{M}_n^1(\mathbb{C})_{sa} \subseteq \mathcal{M}_n(\mathbb{C})_{sa}$  be the smooth function given by*

$$\Psi(U_1, \dots, U_m) = \sum_{j=1}^m w_j^2 U_j^* P_j U_j = \sum_{j=1}^m w_j^2 \Psi_{P_j}(U_j) , \quad \text{for} \quad (U_1, \dots, U_m) \in \mathcal{U}(n)^m .$$

*Then the following conditions are equivalent:*

1. *The differential of  $\Psi$  at  $I := (I_n, \dots, I_n) \in \mathcal{U}(n)^m$  is **surjective***
2.  *$\{P_j : j \in \mathbb{I}_m\}' = \mathbb{C} I_n$ .*

*In this case, the image of  $\Psi$  contains an open neighborhood of  $\Psi(I) = \sum_{j=1}^m w_j^2 P_j$  in  $\mathcal{M}_n^1(\mathbb{C})_{sa}$ , and  $\Psi$  admits smooth (and hence continuous) local cross sections around  $\Psi(I)$ .*

*Proof.* It is clear from its definition that  $\Psi$  is a smooth function. Moreover, under the previous identification  $T_{I_n}\mathcal{U}(n) = \mathcal{M}_n(\mathbb{C})_{ah}$ , and using Eq. (16), we can see that

$$D\Psi_I(X_1, \dots, X_m) = \sum_{j=1}^m w_j^2 [X_j, P_j] \quad , \quad \text{for} \quad (X_1, \dots, X_m) \in \mathcal{M}_n(\mathbb{C})_{ah}^m . \quad (18)$$

The tangent space of  $\mathcal{M}_n^1(\mathbb{C})_{sa}$  is the real vector space  $\mathcal{M}_n^0(\mathbb{C})_{sa}$ , which has a natural inner product given by  $\langle Y, Z \rangle = \text{tr}(YZ)$ . Denote by  $T = D\Psi_I$  and assume that  $T$  is not surjective. Then there exists  $0 \neq Y \in \mathcal{M}_n^0(\mathbb{C})_{sa}$  which is orthogonal to the image of  $T$ . Using Eq. (18) we deduce that, for every  $(X_1, \dots, X_m) \in \mathcal{M}_n(\mathbb{C})_{ah}^m$ , it holds that

$$0 = \langle T(X_1, \dots, X_m), Y \rangle = \sum_{j=1}^m w_j^2 \text{tr}([X_j, P_j]Y) = \sum_{j=1}^m w_j^2 \text{tr}(X_j[P_j, Y]) . \quad (19)$$

Since each  $[P_j, Y] \in \mathcal{M}_n(\mathbb{C})_{ah}$ , we can choose each  $X_j = [P_j, Y]$ , and so Eq. (19) implies that  $[P_j, Y] = 0$  for every  $j \in \mathbb{I}_m$ . In other words, that  $Y \in \{P_j : j \in \mathbb{I}_m\}'$ . On the other hand, since  $0 \neq Y \in \mathcal{M}_n(\mathbb{C})_{sa}$  and  $\text{tr} Y = 0$ , then  $Y \notin \mathbb{C}I_n$ . The converse follows from the previous argument, by taking  $Y \in \{P_j : j \in \mathbb{I}_m\}'$  such that  $0 \neq Y = Y^*$  and  $\text{tr} Y = 0$ .  $\square$

Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. We shall consider on  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  the distance

$$d_P(\mathcal{W}_w, \mathcal{W}'_w) = \max_{i \in \mathbb{I}_m} \|P_{W_i} - P_{W'_i}\|$$

(recall that the weights are fixed), called *punctual*, and the pseudo-distance

$$d_S(\mathcal{W}_w, \mathcal{W}'_w) = \|S_{\mathcal{W}_w} - S_{\mathcal{W}'_w}\| ,$$

called *operatorial*. The problem of finding *local* minimizers for the FF-potential can be stated for anyone of those distances between FF's.

**Corollary 4.2.2.** *Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. Assume that  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  satisfies that  $\{P_{W_j} : j \in \mathbb{I}_m\}' = \mathbb{C}I_n$ . Then  $\mathcal{W}_w$  is a FF and the map*

$$S : \mathcal{B}_{m,n}(\mathbf{d}, w) \rightarrow \mathcal{M}_n^1(\mathbb{C})_{sa} \quad \text{given by} \quad S(\mathcal{V}_w) = S_{\mathcal{V}_w} = \sum_{i \in \mathbb{I}_m} w_i^2 P_{V_i} ,$$

for  $\mathcal{V}_w = (w_i, V_i)_{i \in \mathbb{I}_m} \in \mathcal{B}_{m,n}(\mathbf{d}, w)$ , satisfies that

1. *The image of  $S$  contains an open neighborhood of  $S_{\mathcal{W}_w}$  in  $\mathcal{M}_n^1(\mathbb{C})_{sa}$ .*
2.  *$S$  has  $d_P$ -continuous local cross sections around  $S_{\mathcal{W}_w}$ .*

*Proof.* Notice that the condition  $\{P_{W_j} : j \in \mathbb{I}_m\}' = \mathbb{C}I_n$  implies that  $\mathcal{W}$  is a generating sequence of subspaces. To prove the properties of the map  $S$ , just compose a local cross section for the map  $\Psi$  of Theorem 4.2.1 (which is open in  $I$ ) with the map  $\Phi : \mathcal{U}(n)^m \rightarrow \mathcal{B}_{m,n}(\mathbf{d})$  given by

$$\Phi(U_1, \dots, U_m) = (w_i, U_i(W_i))_{i \in \mathbb{I}_m} = (w_i, R(U_i P_{W_i} U_i^*))_{i \in \mathbb{I}_m} .$$

Observe that  $S \circ \Phi = \Psi$ , so that  $S$  is open in  $\Phi(I) = \mathcal{W}_w$ .  $\square$

**Remark 4.2.3.** Here is an alternative statement of Corollary 4.2.2: Under the same assumptions and notations about  $\mathcal{W}_w$ , it holds that  $S_{\mathcal{W}_w} \in \mathcal{G}l(n)^+$  and, for every sequence  $(S_k)_{k \in \mathbb{N}}$  in  $\mathcal{M}_n^1(\mathbb{C})_{sa}$  such that  $S_k \xrightarrow[k \rightarrow \infty]{} S_{\mathcal{W}_w}$ , there exists a sequence  $(\mathcal{V}_k)_{k \geq k_0}$  in  $\mathcal{S}_{m,n}(\mathbf{d}, w)$  such that

$$d_P(\mathcal{V}_k, \mathcal{W}_w) \xrightarrow[k \rightarrow \infty]{} 0 \quad \text{and} \quad S_{\mathcal{V}_k} = S_k \quad \text{for every} \quad k_0 \leq k \in \mathbb{N}.$$

This formulation of Corollary 4.2.2 generalizes [18, Thm 5.3] to the context of fusion frames with fixed weights.  $\triangle$

It is not clear that a  $d_P$ -local minimizer for the FF-potential on  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  must be a fusion frame, i.e. its frame operator is an invertible operator. The following Lemma shows that this is true.

**Lemma 4.2.4.** *Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. Let  $\mathcal{W}_w$  be a  $d_P$ -local minimizer for the FF-potential in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$ . Then  $S_{\mathcal{W}_w}$  is invertible (equivalently,  $\mathcal{W}_w \in \mathcal{S}_{m,n}(\mathbf{d}, w)$ ).*

*Proof.* Suppose that  $S_{\mathcal{W}_w}$  has nontrivial nullspace  $N(S_{\mathcal{W}_w})$ . If  $x \in N(S_{\mathcal{W}_w})$

$$0 = \langle S_{\mathcal{W}_w} x, x \rangle = \sum_{j \in \mathbb{I}_m} \langle w_j^2 P_{W_j} x, x \rangle = \sum_{j \in \mathbb{I}_m} w_j^2 \|P_{W_j} x\|^2. \quad (20)$$

In other words,  $W_i \subseteq R(S_{\mathcal{W}_w})$  for every  $i \in \mathbb{I}_m$ . Since  $\text{tr}(\mathbf{d}) \geq n > \dim R(S_{\mathcal{W}_w})$ , we deduce that there exists  $i \neq j$  in  $\mathbb{I}_m$  such that  $P_{W_j} P_{W_i} \neq 0$ . Fix that pair  $i, j$ . Fix also  $f \in W_i \setminus W_j^\perp$  and  $g \in N(S_{\mathcal{W}_w})$  two unit vectors. For every  $t \in [0, \pi/2]$ , take the unit vector  $g(t) = \cos t \cdot f + \sin t \cdot g$ .

Let  $V_i = W_i \ominus \text{span}\{f\}$ ,  $W_i(t) = V_i \oplus \text{span}\{g(t)\}$  and  $\mathcal{W}_w(t) \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  the sequence obtained by replacing  $W_i$  by  $W_i(t)$  in  $\mathcal{W}_w$ . As  $g \in W_k^\perp$  for every  $k \in \mathbb{I}_m$ , for every  $t \in (0, \pi/2]$ ,

$$\begin{aligned} \frac{\text{FFP}(\mathcal{W}_w) - \text{FFP}(\mathcal{W}_w(t))}{2} &= \sum_{k=1}^m w_i^2 w_k^2 \left( \text{tr}(P_{W_i} P_{W_k}) - \text{tr}(P_{W_i(t)} P_{W_k}) \right) \\ &= \sum_{k=1}^m w_i^2 w_k^2 \left( \text{tr}(P_{W_i} P_{W_k}) - \text{tr}(g(t)g(t)^* P_{W_k}) - \text{tr}(P_{V_i} P_{W_k}) \right) \\ &= \sum_{k=1}^m w_i^2 w_k^2 \left( \text{tr}(P_{W_i} P_{W_k}) - \cos^2 t \text{tr}(f f^* P_{W_k}) - \text{tr}(P_{V_i} P_{W_k}) \right) \\ &> \sum_{k=1}^m w_i^2 w_k^2 \left( \text{tr}(P_{W_i} P_{W_k}) - \text{tr}(f f^* P_{W_k}) - \text{tr}(P_{V_i} P_{W_k}) \right) = 0, \end{aligned}$$

because  $\text{tr}(f f^* P_{W_j}) = \|P_{W_j} f\|^2 \neq 0$ . Hence  $\text{FFP}(\mathcal{W}_w(t)) < \text{FFP}(\mathcal{W}_w)$  for every  $t \in (0, \pi/2]$ .

Taking  $t \rightarrow 0$ , we have that  $\mathcal{W}_w(t) \xrightarrow{d_P} \mathcal{W}_w$ , and this contradicts the minimality of  $\mathcal{W}_w$ .  $\square$

Given  $S \in \mathcal{M}_n(\mathbb{C})_{sa}$  with  $\sigma(S) = \{\mu_1, \dots, \mu_r\}$ , we denote by  $P_{\mu_k}(S) = P_{N(S - \mu_k I_n)} \in \mathcal{M}_n(\mathbb{C})^+$ , the spectral projection of  $S$  relative to  $\mu_k$ , for  $k \in \mathbb{I}_r$ . These projections satisfy that

1.  $P_{\mu_k}(S) P_{\mu_j}(S) = 0$  if  $k \neq j$ , and  $\sum_{k=1}^p P_{\mu_k}(S) = I_n$  (i.e., they are a system of projectors).

2. For every  $k \in \mathbb{I}_r$ , it holds that  $S P_{\mu_k}(S) = \mu_k P_{\mu_k}(S)$ , so that  $S = \sum_{k=1}^p \mu_k P_{\mu_k}(S)$ .

The following theorem generalizes a similar result given in the paper by Casazza and Fickus [7, Theorem 4], for the case  $w = \mathbb{1}_m$ . Nevertheless, our approach is based on completely different techniques.

**Theorem 4.2.5.** *Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. Let  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  be a local minimizer of the FF-potential with respect to the distance  $d_P$ . If  $\sigma(S_{\mathcal{W}_w}) = \{\mu_1, \dots, \mu_r\}$ , then*

$$P_{\mu_k}(S_{\mathcal{W}_w}) \in C_{\mathcal{W}_w} = \{P_{W_j} : j \in \mathbb{I}_m\}' \quad \text{for every } k \in \mathbb{I}_r. \quad (21)$$

*The same property holds whenever  $\mathcal{W}_w$  is a  $d_P$ -local minimizer in  $\mathcal{B}_{m,n}^1(\mathbf{d})$ .*

*Proof.* Recall from Lemma 4.2.4 that  $\mathcal{W}_w \in \mathcal{S}_{m,n}(\mathbf{d}, w)$ , in other words that  $0 \notin \sigma(S_{\mathcal{W}_w})$ . Consider the set  $\mathcal{Q}$  of finite systems of projectors  $\{Q_k\}_{k \in \mathbb{I}_p}$  such that each  $Q_k \in C_{\mathcal{W}_w}$ . Observe that  $\mathcal{Q}$  is not empty because  $\{I_n\} \in \mathcal{Q}$ . Then  $\mathcal{Q}$  has a maximal element  $\{Q_k\}_{k \in \mathbb{I}_p}$  with respect to the order induced by refinement. Fix  $k \in \mathbb{I}_p$ . For each  $i \in \mathbb{I}_m$  put  $\mathcal{M}_i = W_i \cap R(Q_k)$  and  $\mathcal{N}_i = W_i \cap R(Q_k)^\perp$ . Using that  $Q_k \in C_{\mathcal{W}_w}$ , we get that each  $W_i = \mathcal{M}_i \oplus \mathcal{N}_i$ . Set  $r_i = \dim \mathcal{M}_i$  and  $\mathbf{r} = (r_1, \dots, r_m)$ . Then, the sequence  $\mathcal{W}_{k,w} = (w_i, \mathcal{M}_i)_{i \in \mathbb{I}_m}$  is a FF for  $R(Q_k)$ . We claim that  $\mathcal{W}_{k,w}$  is a local minimizer of the FF-potential in

$$\mathcal{B}(Q_k, \mathbf{r}, w) := \{ \mathcal{V}_w = (w_i, V_i)_{i \in \mathbb{I}_m} \in \mathcal{B}_{m,n}(\mathbf{r}, w) : V_i \subseteq R(Q_k) \text{ for every } i \in \mathbb{I}_m \}.$$

Indeed, given  $\mathcal{V}_w \in \mathcal{B}(Q_k, \mathbf{r}, w)$ , put  $\tilde{\mathcal{V}}_w = (w_i, V_i \oplus \mathcal{N}_i)_{i \in \mathbb{I}_m} \in \mathcal{B}_{m,n}(\mathbf{d}, w)$ . Observe that the map  $\mathcal{V}_w \mapsto \tilde{\mathcal{V}}_w$  preserves the distance  $d_P$ . Moreover, since  $Q_k \in C_{\mathcal{W}_w}$ , then each  $P_{\mathcal{N}_i} = (I_n - Q_k)P_{W_i}$  so that, by Eq (6),  $\text{FFP}(\tilde{\mathcal{V}}_w) = \text{FFP}(\mathcal{V}_w) + \text{FFP}((w_i, \mathcal{N}_i)_{i \in \mathbb{I}_m})$ , and the second summand does not depend on  $\mathcal{V}_w$ . Then, the claim follows from the fact that  $\mathcal{W}_{k,w} = \mathcal{W}_w$ .

Observe that  $S_{\mathcal{W}_w}$  commutes with  $Q_k$ . We now show that  $S_{\mathcal{W}_w} Q_k = \alpha_k Q_k$  for some  $\alpha_k \in \sigma(S_{\mathcal{W}_w})$ . Indeed, by the maximality of  $\{Q_i\}_{i=1}^p$  in  $\mathcal{Q}$ , it follows that there is no non-trivial sub-projection  $Q'$  of  $Q_k$  such that  $Q' \in \{P_{\mathcal{M}_j} : j \in \mathbb{I}_m\}'$ . Then we can apply Corollary 4.2.2 (taking  $\mathcal{H} = R(Q_k)$  and renormalizing the traces) to show that every positive operator (with the correct trace) near  $S_{\mathcal{W}_w} Q_k$  has the form  $S_{\mathcal{V}_w}$  for some  $\mathcal{V}_w \in \mathcal{B}(Q_k, \mathbf{r}, w)$  close to  $\mathcal{W}_k$ . But if  $S_{\mathcal{W}_w} Q_k$  is not a scalar multiple of  $Q_k$ , then we can choose  $S_{\mathcal{V}_w}$  in such a way that

$$\text{FFP}(\mathcal{V}_w) = \text{tr } S_{\mathcal{V}_w}^2 < \text{tr } (S_{\mathcal{W}_w}^2 Q_k) = \text{FFP}(\mathcal{W}_k).$$

But this contradicts the fact that  $\mathcal{W}_k$  is a local minimizer of the FF-potential in  $\mathcal{B}(Q_k, \mathbf{r}, w)$ . Hence  $S_{\mathcal{W}_w} Q_k = \alpha_k Q_k$  and  $Q_k \leq P_{\alpha_k}(S_{\mathcal{W}_w})$ . Using that  $\sum_{k=1}^p Q_k = I_n$ , it is easy to see that each

$$P_{\mu_i}(S_{\mathcal{W}_w}) = \sum_{k \in J_i} Q_k \in C_{\mathcal{W}_w}, \quad \text{where } J_i = \{k \in \mathbb{I}_p : \alpha_k = \mu_i\}. \quad \square$$

**Remark 4.2.6.** Next we give two reinterpretations of Theorem 4.2.5. Under the assumptions and notations of the theorem, the following properties hold:

1. For each  $i \in \mathbb{I}_m$ , there exists an o.n.b.  $\mathcal{B}_i = \{e_j^{(i)} : j \in \mathbb{I}_{d_i}\}$  of  $W_i$ , consisting of eigenvectors of  $S_{\mathcal{W}_w}$ . Indeed, observe that each  $P_{W_i} = \sum_{k \in \mathbb{I}_r} P_{W_i} P_{\mu_k}(S_{\mathcal{W}_w})$  and the fact that, for a fixed  $i \in \mathbb{I}_m$ , the projections  $P_{W_i} P_{\mu_k}(S_{\mathcal{W}_w})$  are pairwise orthogonal.
2. For each  $\mu_k \in \sigma(S_{\mathcal{W}_w})$ , denote by  $\mathcal{M}_{k,i} = N(S_{\mathcal{W}_w} - \mu_k I_n) \cap W_i$ . Then, it holds that the sequence  $\mathcal{W}_k = (w_i, \mathcal{M}_{k,i})_{i \in \mathbb{I}_m}$  is a **tight** FF for  $N(S_{\mathcal{W}_w} - \mu_k I_n)$ . This follows because its frame operator  $S_{\mathcal{W}_k} = P_{\mu_k}(S_{\mathcal{W}_w}) S_{\mathcal{W}_w} = \mu_k P_{\mu_k}(S_{\mathcal{W}_w})$ .  $\triangle$

### 4.3 The eigenvalues of all $d_S$ -minimizers coincide

Recall that, given  $S \in \mathcal{M}_n(\mathbb{C})_{sa}$ , we denote by  $\lambda(S) \in \mathbb{R}^n$  the vector of the  $n$  eigenvalues of  $S$ , counted with multiplicities, in such a way that  $\lambda(S) = \lambda(S)^\downarrow$ .

**Lemma 4.3.1.** *Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. Then*

1.  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  is  $d_P$ -compact.
2. The set  $\Lambda_{m,n}(\mathbf{d}, w) = \{\lambda(S_{\mathcal{W}_w}) : \mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)\}$  is a **convex** and compact subset of  $\mathbb{R}^n$ .
3. The set  $\{S_{\mathcal{W}_w} : \mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)\}$  is compact and closed under unitary conjugation.

*Proof.* Let  $\mathcal{W}_w, \mathcal{W}'_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$ . For  $t \in [0, 1]$  consider  $\lambda = t \lambda(S_{\mathcal{W}_w}) + (1-t) \lambda(S_{\mathcal{W}'_w})$  and notice that  $\lambda = \lambda^\downarrow$ . Therefore, for every admissible  $(m+1)$ -tuple  $(J_0, \dots, J_m) \in LR_r^n(m)$ ,  $1 \leq r \leq n-1$  we have

$$\sum_{j \in J_0} \lambda_j = \sum_{i=1}^m \sum_{j \in J_i} t \lambda_j(S_{\mathcal{W}_w}) + (1-t) \lambda_j(S_{\mathcal{W}'_w}) \leq \sum_{i=1}^m \sum_{j \in J_i} w_j^2 |\{1, \dots, d_j\} \cap J_i| ,$$

since both  $\lambda(S_{\mathcal{W}_w})$  and  $\lambda(S_{\mathcal{W}'_w})$  satisfy Horn-Klyachko's inequalities. Hence, by Theorem 2.3.3, there exists  $\mathcal{V}_w = (w_i, V_i)_{i \in \mathbb{I}_m} \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  such that  $\lambda(S_{\mathcal{V}_w}) = \lambda$ . This shows that  $\lambda \in \Lambda_{m,n}(\mathbf{d}, w)$ , so that  $\Lambda_{m,n}(\mathbf{d}, w)$  is convex. The fact that the set  $\{S_{\mathcal{W}_w} : \mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)\}$  is closed under unitary conjugation is apparent. Finally  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  is  $d_P$ -compact because each Grassmann manifold

$$\mathcal{P}_{d_i}(n) = \{P = P^*P \in \mathcal{M}_n(\mathbb{C})^+ : \text{tr } P = d_i\} = \{UP_iU^* : U \in \mathcal{U}(n)\} = \mathcal{U}(P_i) ,$$

for every fixed  $P_i \in \mathcal{P}_{d_i}(n)$ , is compact. This follows because  $\mathcal{U}(n)$  is compact. By continuity, the other sets involved are also compact.  $\square$

**Theorem 4.3.2.** *Let  $(\mathbf{d}, w) \in \mathbb{N}^m \times \mathbb{R}_{>0}^m$  be a normalized pair. Then,*

1. The spectra (with multiplicities) of all the frame operators of global minimizers of the FF-potential in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  coincide.
2. The local minimizers of the FF-potential in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  with respect to the pseudo-distance  $d_S$  lie in  $\mathcal{S}_{m,n}(\mathbf{d}, w)$  and are also global minimizers.

*Proof.* Let  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  and notice that  $\text{FFP}(\mathcal{W}_w) = \|\lambda(S_{\mathcal{W}_w})\|^2$ . Since  $\Lambda_{m,n}(\mathbf{d}, w)$  is a compact convex set, then there exists a **unique**  $\lambda_0 \in \Lambda_{m,n}(\mathbf{d}, w)$  which minimizes the euclidean norm on  $\Lambda_{m,n}(\mathbf{d}, w)$ . Hence, if  $\mathcal{W}_w$  is a global minimizer of the FF-potential in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$ , we can conclude that  $\|\lambda(S_{\mathcal{W}_w})\|^2 \leq \|\lambda_0\|^2$ , which implies that  $\lambda(S_{\mathcal{W}_w}) = \lambda_0$ .

Observe that the map  $\sigma : \mathcal{B}_{m,n}(\mathbf{d}, w) \rightarrow \Lambda_{m,n}(\mathbf{d}, w)$  given by  $\sigma(\mathcal{W}_w) = \lambda(S_{\mathcal{W}_w})$  is continuous with respect to the pseudo-distance  $d_S$  of  $\mathcal{B}_{m,n}(\mathbf{d}, w)$ . Moreover,  $\sigma$  is an open map. Indeed, fix  $\mathcal{W}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$ ,  $\lambda = \lambda(S_{\mathcal{W}_w}) \in \Lambda_{m,n}(\mathbf{d}, w)$  and take  $\mu \in \Lambda_{m,n}(\mathbf{d}, w)$  close to  $\lambda$ . Let  $U \in \mathcal{U}(n)$  such that  $S_{\mathcal{W}_w} = U \text{diag}(\lambda) U^*$ . By Lemma 4.3.1, there exists  $\mathcal{V}_w \in \mathcal{B}_{m,n}(\mathbf{d}, w)$  such that  $S_{\mathcal{V}_w} = U \text{diag}(\mu) U^*$ . Now observe that  $d_S(\mathcal{W}_w, \mathcal{V}_w) = \|S_{\mathcal{W}_w} - S_{\mathcal{V}_w}\| = \|\lambda - \mu\|_\infty$ .

Therefore, if  $\mathcal{W}_w$  is a  $d_S$ -local minimizer of the FF-potential in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$ , then  $\lambda(S_{\mathcal{W}_w})$  is a local minimum for the euclidean norm in the set  $\Lambda_{m,n}(\mathbf{d}, w) \subseteq \mathbb{R}^n$ . By a standard computation, the convexity of  $\Lambda_{m,n}(\mathbf{d}, w)$  implies that  $\lambda(S_{\mathcal{W}_w})$  must be the global minimizer  $\lambda_0$ , and therefore  $\mathcal{W}_w$  is a global minimizer in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$ .  $\square$



**Conjecture 4.3.3.** Local minimizers of the frame potential in  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  (resp.  $\mathcal{S}_{m,n}^1(\mathbf{d})$ ) with respect to punctual distance  $d_P$  are also global minimizers.

In some particular cases (i.e. for particular choices of the parameters  $n, m, \mathbf{d}$  and  $w$ ), there is an affirmative answer for this conjecture (see [7, Theorem 5]).

**Remark 4.3.4.** All the previous results remain true if one replaces  $\mathcal{B}_{m,n}(\mathbf{d}, w)$  by  $\mathcal{B}_{m,n}^1(\mathbf{d})$  (i.e., minimizing without fixing the sequence of weights). We present some of the new statements without proofs, since they are based on techniques that are similar to those already developed.

1. As in the proof Lemma 4.3.1, Horn-Klyachko's compatibility inequalities (8) show that

$$\Lambda_{m,n}^1(\mathbf{d}) = \{\lambda(S_{\mathcal{W}_w}) : \mathcal{W}_w \in \mathcal{B}_{m,n}^1(\mathbf{d})\} \quad \text{is compact and convex.}$$

2. This fact implies that the (ordered) spectra of the frame operators of global minimizers of the FF-potential in  $\mathcal{B}_{m,n}^1(\mathbf{d})$  coincide.
3. Finally, the argument of the proof of Theorem 4.3.2 can be adapted to yield that local minimizers of the FF-potential in  $\mathcal{B}_{m,n}^1(\mathbf{d})$ , with respect to the operator distance  $d_S$ , are also global.  $\triangle$

As in the case of fixed weights (Lemma 4.2.4), the global minimizers for the FF-potential in  $\mathcal{B}_{m,n}^1(\mathbf{d})$  are fusion frames.

**Proposition 4.3.5.** Let  $\mathcal{W}_w$  a  $d_S$ -local (and hence global) minimizer for the FF-potential in  $\mathcal{B}_{m,n}^1(\mathbf{d})$ . Then its frame operator  $S_{\mathcal{W}_w} \in \mathcal{GL}(n)^+$ . In other words,  $\mathcal{W}_w \in \mathcal{S}_{m,n}^1(\mathbf{d})$ .

*Proof.* Let  $J = \{i \in \mathbb{I}_m : w_i \neq 0\}$ , and  $k = \sum_{i \in J} d_i$ . Note that, if  $k \geq n$ , we can apply Lemma 4.2.4 (fixing the weight  $w_J$ ) and we are done.

We assume that  $k < n$ , and will obtain a contradiction. Without loss of generality, we can suppose that  $J = \mathbb{I}_r$ . It follows immediately that  $W_i \perp W_j$  for  $1 \leq i \neq j \leq r$  and  $w_i^2 = \frac{1}{k}$  for every  $1 \leq i \leq r$ . Thus,  $\text{FFP}(\mathcal{W}_w) = \frac{1}{k}$ . Moreover, if  $d = d_{r+1}$ , then  $k + d > n$ . Otherwise, if we take a subspace  $W_{r+1} \subseteq (\bigcup_{i \in \mathbb{I}_r} W_i)^\perp$  with  $\dim W_{r+1} = d$  and we set  $w_i^2 = \frac{1}{k+d}$ , for  $i \in \mathbb{I}_{r+1}$ , then we get a BSS in  $\mathcal{B}_{m,n}^1(\mathbf{d})$  with FF-potential  $\frac{1}{k+d} < \frac{1}{k}$ .

Therefore, we can construct  $\mathcal{V}_v = (v_j, V_j)_{j \in \mathbb{I}_{r+1}} \in \mathcal{B}_{m,n}^1(\mathbf{d})$  in the following way:

$$v_j^2 = \begin{cases} a & \text{if } 1 \leq j \leq r \\ b & \text{if } j = r+1 \end{cases} \quad \text{and} \quad V_j = \begin{cases} W_j & \text{if } 1 \leq j \leq r \\ [\bigoplus_{i=1}^r W_i]^\perp \oplus T & \text{if } j = r+1 \end{cases},$$

where  $T \subseteq \bigoplus_{i=1}^r W_i$  is a subspace with  $\dim T = d + k - n$  (so that  $\dim V_{r+1} = d$ ), and  $ka + db = 1$ . It is easy to see, by taking an orthonormal basis of each subspace  $V_i$ , that

$$f(a) = \text{FFP}(\mathcal{V}_v) = ka^2 + db^2 + 2(d + k - n)ab \quad \text{with} \quad ka + db = 1, \quad a \in [0, \frac{1}{k}] \quad (22)$$

Easy computations show that  $f'(\frac{1}{k}) = \frac{2(n-k)}{d} > 0$ . Since  $f(\frac{1}{k}) = \frac{1}{k}$ , there exist pairs  $(a, b)$  such that the FF-potential is lower than  $\frac{1}{k}$ , which contradicts the minimality of  $\mathcal{W}_w$ .  $\square$

Next, we summarize the facts described in Remark 4.3.4 and Proposition 4.3.5.

**Theorem 4.3.6.** Let  $\mathbf{d} \in \mathbb{N}^m$ . Then, local minimizers of the frame potential in  $\mathcal{B}_{m,n}^1(\mathbf{d})$  with respect to the pseudo-distance  $d_S$  lie in  $\mathcal{S}_{m,n}^1(\mathbf{d})$  and are also global minimizers. Moreover, the spectra of the frame operators of these local minimizers coincide.  $\square$

## 5 Minimization for fixed subspaces

In this section we shall characterize the sequences of weights which minimize the potential of a fixed sequence of subspaces. The main tools are some results about Hadamard indexes of [12], which we shall state in some detail in the Appendix. Recall that, for  $A, B \in \mathcal{M}_n(\mathbb{C})$ , their Hadamard product is the matrix  $A \circ B = (A_{ij} B_{ij})_{i,j \in \mathbb{I}_n} \in \mathcal{M}_n(\mathbb{C})$ .

### 5.1 Minimal weights

In this section we fix  $m \in \mathbb{N}$ , a Hilbert space  $\mathcal{H}$  with  $\dim \mathcal{H} = n$ , and  $\mathbf{d} \in \mathbb{N}^m$  with  $M = \text{tr } \mathbf{d} \geq n$ . We also fix a sequence  $\mathcal{W} = \{W_i\}_{i \in \mathbb{I}_m}$  of subspaces which spans  $\mathcal{H}$ , such that each  $\dim W_i = d_i$ . Our aim is to minimize the FF-potential over all sequences  $w \in \mathbb{R}_+^m$  such that  $\mathcal{W}_w \in \mathcal{S}_{m,n}^1(\mathbf{d})$ .

Recall that the Benedetto-Fickus FF-potential of  $\mathcal{W}_w$  is given by:

$$\text{FFP}(\mathcal{W}_w) = \sum_{i,j=1}^m w_i^2 w_j^2 \text{tr}(P_{W_i} P_{W_j}) = \text{tr } S_{\mathcal{W}_w}^2. \quad (23)$$

**Definition 5.1.1.** Let  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m} \in \mathcal{S}_{m,n}^1(\mathbf{d})$ . We denote by

$$B = B_{\mathcal{W}_w} \in \mathcal{M}_m(\mathbb{R}) \quad \text{the matrix given by} \quad B_{ij} = w_i^2 w_j^2 \text{tr}(P_{W_i} P_{W_j}). \quad \triangle$$

**Lemma 5.1.2.** Let  $\mathcal{W}_w = (w_i, W_i)_{i \in \mathbb{I}_m}$  be a FF for  $\mathcal{H}$ . Then,  $B_{\mathcal{W}_w} \in \mathcal{M}_m(\mathbb{R})^+$ .

*Proof.*  $B_{\mathcal{W}_w} \in \mathcal{M}_m(\mathbb{R})^+$  because it is the Gram matrix for the vectors  $\{w_i^2 P_{W_i}\}_{i \in \mathbb{I}_m}$  in the euclidean space  $\mathcal{M}_n(\mathbb{C})$  with the inner product defined as  $\langle X, Y \rangle = \text{tr } Y^* X$ .  $\square$

**Notations 5.1.3.** We shall fix some notations and assumptions:

1. We begin with a fixed normalized sequence of weights, in the sense that

$$\mathbf{w} = (w_i)_{i \in \mathbb{I}_m} \quad \text{is given by} \quad w_i = d_i^{-\frac{1}{2}} \quad \text{for every} \quad i \in \mathbb{I}_m.$$

Observe that the condition  $w_i = d_i^{-\frac{1}{2}}$  means that each “vector”  $w_i P_{W_i}$  of  $\mathcal{W}_w$  has size  $\|w_i P_{W_i}\|_2 = 1$ . This justifies the word “normalized” of  $\mathcal{W}_w$ .

2. Given a sequence of weights  $\mathbf{a} = (a_i)_{i \in \mathbb{I}_m} \in \mathbb{R}_+^m$ , we denote by  $\mathbf{a} \cdot \mathcal{W}_w$  the Bessel sequence of subspaces  $\mathbf{a} \cdot \mathcal{W}_w = (a_i w_i W_i)_{i \in \mathbb{I}_m}$ .
3. In  $\mathbf{a} \in \mathbb{R}_+^m$ , then  $\text{tr}(S_{\mathbf{a} \cdot \mathcal{W}_w}) = \sum_{i \in \mathbb{I}_m} a_i^2 w_i^2 d_i$ . Therefore, as we start with normalized weights,

$$\mathbf{a} \cdot \mathcal{W}_w \in \mathcal{S}_{m,n}^1(\mathbf{d}) \iff \sum_{i \in \mathbb{I}_m} a_i^2 w_i^2 d_i = \sum_{i \in \mathbb{I}_m} a_i^2 = 1 \iff \|\mathbf{a}\| = 1. \quad (24)$$

4. Let  $A = A_{\mathcal{W}_w} \in \mathcal{M}_m(\mathbb{R})$  the matrix given by  $A_{ij} = (B_{\mathcal{W}_w})_{ij}^{\frac{1}{2}} = w_i w_j \text{tr}(P_{W_i} P_{W_j})^{\frac{1}{2}}$ , for  $i, j \in \mathbb{I}_m$ . Observe that  $A$  is selfadjoint, but possibly  $A \notin \mathcal{M}_m(\mathbb{R})^+$ . On the other hand,

$$\begin{aligned} \text{FFP}(\mathcal{W}_w) &= \sum_{i,j=1}^m w_i^2 w_j^2 \text{tr}(P_{W_i} P_{W_j}) = \|A_{\mathcal{W}_w}\|_2^2 \quad \text{and also} \\ \text{FFP}(\mathbf{a} \cdot \mathcal{W}_w) &= \sum_{i,j=1}^m a_i^2 a_j^2 w_i^2 w_j^2 \text{tr}(P_{W_i} P_{W_j}) = \|\mathbf{a} \mathbf{a}^* \circ A_{\mathcal{W}_w}\|_2^2. \end{aligned} \quad (25)$$

5. Using the previous identities, we can now define the main notion of this section:

$$I(\mathcal{W}_w) := \min_{\|\mathbf{a}\|=1} \text{FFP}(\mathbf{a} \cdot \mathcal{W}_w) = \min_{\|\mathbf{a}\|=1} \|\mathbf{a}\mathbf{a}^* \circ A_{\mathcal{W}_w}\|_2^2. \quad \triangle$$

In order to compute  $I(\mathcal{W}_w)$  as well as to describe the set of weights  $\mathbf{a} \in \mathbb{R}_{\geq 0}^m$ , with  $\|\mathbf{a}\| = 1$  for which  $I(\mathcal{W}_w) = \text{FFP}(\mathbf{a} \cdot \mathcal{W}_w)$ , the main tools are some results about Hadamard indexes of [12], which we shall state in some detail in the Appendix. Here we just give the basic definitions.

**Definition 5.1.4.** Let  $B \in \mathcal{M}_m(\mathbb{C})^+$ . The **minimal**-Hadamard index of  $G$  is

$$I(B) = \max \{ \lambda \geq 0 : B \circ C \geq \lambda C \quad \text{for every } C \in \mathcal{M}_m(\mathbb{C})^+ \}.$$

For  $A \in \mathcal{M}_m(\mathbb{C})_{sa}$ , we define the spectral and the  $\|\cdot\|_2$  Hadamard indexes:

$$I_{sp}(A) = \min_{\|x\|=1} \|A \circ x x^*\|_{sp} \quad \text{and} \quad I_2(A) = \min_{\|x\|=1} \|A \circ x x^*\|_2.$$

For a matrix  $G \in \mathcal{M}_n(\mathbb{C})$ , we write  $0 \leq G$  if all entries  $G_{ij} \geq 0$ . Given  $J \subseteq \mathbb{I}_m$  with  $|J| = k$  we denote by  $G_J \in \mathcal{M}_k(\mathbb{C})$  the submatrix of  $G$  given by  $G_J = (G_{ij})_{i,j \in J}$ . Similarly, if  $x \in \mathbb{C}^n$ , we write  $x \geq 0$  if  $x \in \mathbb{R}_+^n$  and  $x_J = (x_j)_{j \in J} \in \mathbb{C}^k$ .  $\triangle$

From the previous definitions and Eq. (25), we get the fundamental equality:  $I(\mathcal{W}_w) = I_2(A_{\mathcal{W}_w})$ . Now it is clear why the results of the Appendix can be useful for computing  $I(\mathcal{W}_w)$ .

**Remark 5.1.5.** Let  $\mathbf{a} \in \mathbb{R}^m$ . Then

$$\text{FFP}(\mathbf{a} \cdot \mathcal{W}_w) = \sum_{i,j=1}^m a_i^2 a_j^2 w_i^2 w_j^2 \text{tr}(P_i P_j) = \langle (B \circ \mathbf{a}\mathbf{a}^*) \mathbf{a}, \mathbf{a} \rangle = \langle B(\mathbf{a} \circ \mathbf{a}^*), (\mathbf{a} \circ \mathbf{a}^*) \rangle \quad (26)$$

where, as before,  $B = B_{\mathcal{W}_w} \in \mathcal{M}_m(\mathbb{R})^+$ . Moreover, by Eq. (32) in Proposition A.1.2,

$$I(\mathcal{W}_w) = \min_{\|\mathbf{a}\|=1} \|A_{\mathcal{W}_w} \circ \mathbf{a}\mathbf{a}^*\|_2^2 = \min_{\|\mathbf{a}\|=1} \|B \circ \mathbf{a}\mathbf{a}^*\| = I_{sp}(B). \quad (27)$$

This identity is useful because  $0 \leq B \in \mathcal{M}_m(\mathbb{R})^+$  and its spectral index is easier to compute. Indeed, observe that  $\text{tr } P_{W_i} P_{W_j} \geq 0$  for every  $i, j \in \mathbb{I}_m$ , since each  $P_{W_i} \in \mathcal{M}_n(\mathbb{C})^+$ .  $\triangle$

## 5.2 Critical Points and local minimizers.

In what follows, we shall use all the assumptions and notations of the previous subsection, but we need the following extra notations:

1. Given  $\mathbf{a} \in \mathbb{R}^m$ , we write  $z = \mathbf{a} \circ \mathbf{a} = (a_1^2, \dots, a_m^2)$  and  $J = \text{supp}\{\mathbf{a}\} = \{j \in \mathbb{I}_n : a_j \neq 0\}$ .
2. Let  $L : \mathbb{R}^m \rightarrow \mathbb{R}^+$  be given by  $L(\mathbf{a}) = \text{FFP}(\mathbf{a} \cdot \mathcal{W}_w) = \|A_{\mathcal{W}_w} \circ \mathbf{a}\mathbf{a}^*\|_2^2 = \langle B_{\mathcal{W}_w} z, z \rangle$ .
3. We consider the affine manifold  $\Delta_0 = \{x \in \mathbb{R}^m : \text{tr } x = 0\}$  and the compact convex simplex  $\Delta = \{x \in \Delta_0 : x \geq 0\}$ .
4.  $S^{m-1} = \{x \in \mathbb{R}^m : \|x\| = 1\}$  is the unit sphere of  $\mathbb{R}^m$ .

**Lemma 5.2.1.** Let  $0 \leq \mathbf{a} \in S^{m-1}$ . Denote  $B = B_{\mathcal{W}_w}$ . The following conditions are equivalent:

1.  $\mathbf{a}$  is a critical point for the map  $L$  restricted to the sphere  $S^{m-1}$ .

2.  $B_J z_J = I(B_J) \mathbf{1}_J$ , where  $J = \text{supp}\{\mathbf{a}\}$ .

*Proof.* Observe that  $L(\mathbf{a}) = \langle Bz, z \rangle = \sum_{i,j=1}^m b_{ij} a_i^2 a_j^2$ . Since  $B = B^* = B^T$ , we have that

$$0 = \nabla L(\mathbf{a}) = 4 \left( a_1 \sum_{j=1}^m b_{1j} a_j^2, \dots, a_m \sum_{j=1}^m b_{mj} a_j^2 \right) = 4 Bz \circ \mathbf{a}.$$

The tangent space of  $S^{m-1}$  at  $\mathbf{a}$  is  $\{\mathbf{a}\}^\perp$ . Hence,  $\mathbf{a}$  is a critical point for  $S^{m-1}$  if and only if

$$0 = \langle \nabla L(\mathbf{a}), y \rangle = 4 \langle Bz \circ \mathbf{a}, y \rangle \quad \text{for every } y \in \{\mathbf{a}\}^\perp \iff Bz \circ \mathbf{a} \in \text{span}\{\mathbf{a}\}.$$

This is clearly equivalent to the equation  $B_J z_J = \lambda \mathbf{1}_J$ , for some  $\lambda \in \mathbb{R}$ . In this case, since  $0 \leq B_{ij}$  with  $0 < B_{ii}$  for every  $i \in \mathbb{I}_n$  and  $0 \leq z$ , we can conclude that  $\lambda > 0$ . Moreover, by Proposition A.1.2 applied to the matrix  $B_J$ , we have that

$$B_J z_J = \lambda \mathbf{1}_J \implies B_J \frac{z_J}{\lambda} = \mathbf{1}_J \implies I(B_J)^{-1} = \text{tr} \frac{z_J}{\lambda} = \langle \mathbf{1}_J, \frac{z_J}{\lambda} \rangle = \lambda^{-1} \langle \mathbf{1}, z \rangle = \lambda^{-1}.$$

Therefore  $\lambda = I(B_J)$  and  $B_J z_J = I(B_J) \mathbf{1}_J$ . □

**Theorem 5.2.2.** *Let  $0 \leq \mathbf{a} \in S^{m-1}$  such that  $B_J z_J = I(B_J) \mathbf{1}_J$ , i.e.,  $\mathbf{a}$  is a critical point of  $L$  restricted to  $S^{m-1}$ . Then, the following conditions are equivalent:*

1.  $I(B_J) = I_{sp}(B) = I(\mathcal{W}_w)$ .
2.  $\mathbf{a}$  is a global minimum of  $L$  restricted to  $S^{m-1}$ .
3.  $\mathbf{a}$  is a local minimum of  $L$  restricted to  $S^{m-1}$ .
4.  $Bz \geq I(B_J) \mathbf{1}$ . In other words, that  $(Bz)_j \geq I(B_J)$  for every  $j \notin J$ .

*Proof.* Denote  $A = A_{\mathcal{W}_w}$ . Recall that  $I_2(A)^2 = I_{sp}(B)$ , by Eq. (27). By Lemma 5.2.1,

$$\|A \circ \mathbf{a} \mathbf{a}^*\|_2^2 = \langle Bz, z \rangle = I(B_J) \langle \mathbf{1}, z \rangle = I(B_J).$$

This gives the equivalence  $1 \leftrightarrow 2$ . Observe that, if  $\mathbf{b} \in S^{m-1}$ , then  $w = \mathbf{b} \circ \mathbf{b} \in \Delta$ . For each  $w \in \Delta$ , consider the line  $\gamma_w : [0, 1] \rightarrow \Delta$  joining  $z$  and  $w$ , given by the formula  $\gamma_w(t) = (1-t)z + tw$ , for every  $t \in [0, 1]$ . Consider the map  $\rho_w : [0, 1] \rightarrow \mathbb{R}^+$  given by

$$\rho_w(t) = \langle B\gamma_w(t), \gamma_w(t) \rangle = (1-t)^2 \langle Bz, z \rangle + t^2 \langle Bw, w \rangle + 2t(1-t) \langle Bz, w \rangle, \quad (28)$$

for every  $t \in [0, 1]$ . Since  $\langle Bz, z \rangle = I(B_J)$ , the derivative of  $\rho_w$  evaluated at zero is

$$\dot{\rho}_w(0) = -2 \langle Bz, z \rangle + 2 \langle Bz, w \rangle = 2 \left( \langle Bz, w \rangle - I(B_J) \right).$$

On the other hand, for every  $t \in \mathbb{R}$ ,

$$\ddot{\rho}_w(t) = 2 \langle Bz, z \rangle + 2 \langle Bw, w \rangle - 4 \langle Bz, w \rangle = 2 \langle B(z-w), z-w \rangle \geq 0. \quad (29)$$

Since  $\rho_w$  is a second degree polynomial, its leading coefficient is  $\frac{1}{2} \ddot{\rho}_w(t) \geq 0$ . Suppose now that  $Bz \geq I(B_J)\mathbf{1}$ . Using that  $w \in \Delta$ , we get that

$$\langle Bz, w \rangle \geq I(B_J) \langle \mathbf{1}, w \rangle = I(B_J) \implies \dot{\rho}_w(0) \geq 0 \implies \dot{\rho}_w(t) \geq 0 \quad \text{for every } t \geq 0.$$

Therefore  $\rho_w(1) \geq \rho_w(0)$ . In other words, we have proved that  $\langle Bw, w \rangle \geq \langle Bz, z \rangle$  for every  $w \in \Delta$ . This implies that  $\mathbf{a}$  is a global minimum of  $L$  restricted to  $S^{m-1}$ .

Suppose now that  $\mathbf{a} \in S^{m-1}$ ,  $z = \mathbf{a} \circ \mathbf{a}$ , and that there exists  $k \in \mathbb{I}_m$  such that  $(Bz)_k < I(B_J)$ . Observe that  $\mathbf{a} \neq e_k$ , because  $(Be_k)_k = b_{kk} = I(B_{\{k\}})$ .

Let  $w = e_k \in \Delta$ , and consider the curves  $\gamma_w$  and  $\rho_w$  defined before. By the previous computations, we have that  $\dot{\rho}_w(0) = 2((Bz)_k - I(B_J)) < 0$ . Therefore, for every  $t > 0$  small enough, we have that  $\gamma_w(t) \in \Delta$  and  $\langle B\gamma_w(t), \gamma_w(t) \rangle < \langle Bz, z \rangle$ . Taking the vectors  $\mathbf{a}(t) = \text{sgn}(\mathbf{a}) \gamma_w(t)^{\frac{1}{2}} \in S^{m-1}$ , we conclude that  $\mathbf{a}$  fails to be a local minimum.  $\square$

**Remark 5.2.3.** Given  $\mathbf{a} \in S^{m-1}$ , then  $z = \mathbf{a} \circ \mathbf{a} \in \Delta$  and  $L(\mathbf{a}) = \langle Bz, z \rangle$ . Therefore

$$I(\mathcal{W}_w) = I_2(A) = \min_{\mathbf{a} \in S^{m-1}} \|A \circ \mathbf{a}\mathbf{a}^*\|_2^2 = \min_{0 \leq \mathbf{a} \in S^{m-1}} \|A \circ \mathbf{a}\mathbf{a}^*\|_2^2 = \min_{z \in \Delta} \langle Bz, z \rangle,$$

since every  $z \in \Delta$  produces a unit vector  $0 \leq \mathbf{a} = z^{\frac{1}{2}} \in S^{m-1}$ . Then in order to get the unit vectors  $\mathbf{a}$  which attain this minimum, it suffices to characterize the sets

$$\mathcal{S}(\mathcal{W}_w) = \arg \min_{z \in \Delta} \{\langle Bz, z \rangle\} \quad \text{and} \quad J(\mathcal{W}_w) = \{J \subseteq \mathbb{I}_m : J = \text{supp}\{z\} \text{ for some } z \in \mathcal{S}(\mathcal{W}_w)\}.$$

If  $\mathbb{I}_m \notin J(\mathcal{W}_w)$ , it is possible to obtain minimizers  $\mathbf{a} \cdot \mathcal{W}_w$  which are not fusion frames, because  $S_{\mathbf{a} \cdot \mathcal{W}_w} \notin \mathcal{GL}(n)^+$  (see Example 5.2.7). Still, if  $I(\mathcal{W}_w) < \frac{\sqrt{1+n}}{n}$  then  $S_{\mathbf{a} \cdot \mathcal{W}_w} \in \mathcal{GL}(n)^+$  for any minimizer  $\mathbf{a}$ , since in such case Proposition 2.2.3 implies that  $\|I - n S_{\mathbf{a} \cdot \mathcal{W}_w}\| < 1$ . Otherwise, the characterization of the set  $J(\mathcal{W}_w)$  is useful in order to discern if there are optimal sequences of weights  $\mathbf{a}$  such that  $\mathbf{a} \cdot \mathcal{W}_w$  remains being a FF. Item 4 of Theorem 5.2.2 gives a description of the elements of  $\mathcal{S}(\mathcal{W}_w)$ . But its proof gives more information:  $\triangle$

**Corollary 5.2.4.** Consider  $\mathcal{W}_w \in \mathcal{S}_{m,n}(\mathbf{d})$ ,  $A = A_{\mathcal{W}_w}$  and  $B = B_{\mathcal{W}_w}$  as before. Then

1. The set  $\mathcal{S}(\mathcal{W}_w) = \arg \min_{z \in \Delta} \{\langle Bz, z \rangle\}$  is convex. Moreover, for any point  $z_0 \in \mathcal{S}(\mathcal{W}_w)$ ,

$$\mathcal{S}(\mathcal{W}_w) = (z_0 + N(B)) \cap \Delta.$$

2.  $J(\mathcal{W}_w)$  is closed under taking unions, so that  $J_{\mathcal{W}_w} = \bigcup J(\mathcal{W}_w) = \bigcup_{z \in \mathcal{S}(\mathcal{W}_w)} \text{supp}\{z\}$  is an element of  $J(\mathcal{W}_w)$ , and there exists  $z_1 \in \mathcal{S}(\mathcal{W}_w)$  with maximal support.

*Proof.* 1. Let  $z, w \in \Delta$ , and consider the function, defined in Eq. (28):

$$\rho_{z,w}(t) = \langle B((1-t)z + tw), (1-t)z + tw \rangle, \quad t \in \mathbb{R}.$$

Suppose now that  $w, z \in \mathcal{S}(\mathcal{W}_w)$  and  $w \neq z$ . Using that  $\rho_{z,w}$  is of second degree, the equality  $\ddot{\rho}_{z,w}(t) = 2 \langle B(z-w), z-w \rangle \geq 0$  given by Eq. (29), and the fact  $\rho_{z,w}(t) \geq 0$  for every  $t \in \mathbb{R}$ , we can conclude that

$$\rho_{z,w} \text{ is constant} \iff \ddot{\rho}_{z,w}(t) = 0 \iff z - w \in N(B).$$

On the other hand, we have that  $\rho_{z,w}(1) = \rho_{z,w}(0) = \min_{t \in [0,1]} \rho_{z,w}(t)$ . This implies that the map  $\rho_{z,w}$  is constant, so that  $\gamma(t) \in \mathcal{S}(\mathcal{W}_w)$  for every  $t \in [0,1]$ , and  $z - w \in N(B)$ . The proof of the fact that  $(z_0 + N(B)) \cap \Delta \subseteq \mathcal{S}(\mathcal{W}_w)$  for every  $z_0 \in \mathcal{S}(\mathcal{W}_w)$  is similar.

2. Let  $z$  and  $w$  in  $\mathcal{S}(\mathcal{W}_w)$ , with supports  $J_1$  and  $J_2$  respectively. Then, since the entire line  $tz + (1-t)w \in \mathcal{S}(\mathcal{W}_w)$  ( $t \in [0, 1]$ ), if we take  $u = tz + (1-t)w$  for any  $t \in (0, 1)$ , it is easy to see that  $u \in \mathcal{S}(\mathcal{W}_w)$  and  $\text{supp}\{u\} = J_1 \cup J_2$ . Since  $J(\mathcal{W}_w)$  is finite, also the set

$$J_{\mathcal{W}_w} = \bigcup_{z \in \mathcal{S}(\mathcal{W}_w)} \text{supp}\{z\} \in J(\mathcal{W}_w) .$$

Hence  $J_{\mathcal{W}_w}$  is the support of some  $z_1 \in \mathcal{S}(\mathcal{W}_w)$ .  $\square$

**Corollary 5.2.5.** Let  $B = B_{\mathcal{W}_w}$  and  $A = A_{\mathcal{W}_w}$  as before. Assume that there exists  $v \in \mathbb{R}^m$  such that  $v \geq 0$  and  $Bv = \mathbb{1}$ . Denote  $\mathbf{a} = (\text{tr } v)^{-\frac{1}{2}} (v_1^{\frac{1}{2}}, \dots, v_m^{\frac{1}{2}})$ . Then

$$\|\mathbf{a}\| = 1 \quad \text{and} \quad (\text{tr } v)^{-1} = \text{FFP}(\mathbf{a} \cdot \mathcal{W}_w) = I_2(A)^2 = I(\mathcal{W}_w) . \quad (30)$$

*Proof.* The fact that  $v \geq 0$  and  $Bv = \mathbb{1}$  implies, by Propositions A.1.2 and A.1.3, that

$$I_2(A)^2 = I_{sp}(B) = I(B) = I(B_J) = (\text{tr } v)^{-1} ,$$

where  $J = \text{supp}\{v\} = \text{supp}\{\mathbf{a}\}$ . Since  $z = \frac{v}{\text{tr } v} \in \Delta$ , then  $\mathbf{a} = z^{\frac{1}{2}} \in S^{m-1}$  and  $Bz = I(B_J)\mathbb{1}$ . Hence, by Theorem 5.2.2, we have that  $z \in \mathcal{S}(\mathcal{W}_w)$  and  $\mathbf{a} \in S^{m-1}$  satisfies Eq. (30).  $\square$

**Remark 5.2.6.** The results of this section seems to be unknown still for the case of vector frames. In this case our restrictions translate to the following: Let  $\mathcal{F} = (f_i)_{i \in \mathbb{I}_m}$  be a frame for  $\mathcal{H}$  such that each  $\|f_i\| = 1$  (i.e.,  $d_i = 1 \implies w_i = d_i^{-\frac{1}{2}} = 1$ ). For  $\mathbf{a} \in \mathbb{R}^m$ , we consider the sequence  $\mathbf{a} \cdot \mathcal{F} = (a_i f_i)_{i \in \mathbb{I}_m}$ , and we define  $I(\mathcal{F}) = \min_{\|\mathbf{a}\|=1} \text{FP}(\mathbf{a} \cdot \mathcal{F})$ . Then, all the results of the section remain true if one consider the matrices

$$A_{\mathcal{F}} = \left( |\langle f_j, f_i \rangle| \right)_{i,j \in \mathbb{I}_m} \in \mathcal{M}_m(\mathbb{C})_{sa} \quad \text{and} \quad B_{\mathcal{F}} = \left( |\langle f_j, f_i \rangle|^2 \right)_{i,j \in \mathbb{I}_m} \in \mathcal{M}_m(\mathbb{C})^+ .$$

Some proofs are slightly easier in this case, because  $I(\mathcal{F}) = I_2(A_{\mathcal{F}}) = I_2(G_{\mathcal{F}})$ , where  $G_{\mathcal{F}}$  is the Gramm matrix of  $\mathcal{F}$ :  $G_{\mathcal{F}} = (\langle f_j, f_i \rangle)_{i,j \in \mathbb{I}_m} \in \mathcal{M}_m(\mathbb{C})^+$ . Observe that the diagonal entries of the three matrices involved are equal to 1.  $\triangle$

**Example 5.2.7.** Let  $B = \frac{1}{4} \begin{pmatrix} 4 & 1 & 3 \\ 1 & 4 & 2 \\ 3 & 2 & 4 \end{pmatrix}$ . Since  $A = (B_{ij}^{\frac{1}{2}})_{i,j \in \mathbb{I}_3} \in \mathcal{G}l(3)^+$ , we deduce that  $A$  is the Gramm matrix of a Riesz basis  $\mathcal{F}$  of  $\mathbb{C}^3$ . Let  $v = (\frac{4}{5}, \frac{4}{5}, 0) \geq 0$ . Observe that

$$Bv = \mathbb{1}_3 \implies \quad (\text{by Corollary 5.2.5}) \quad z = (\text{tr } v)^{-1} v = \left( \frac{1}{2}, \frac{1}{2}, 0 \right) \in \mathcal{S}(\mathcal{F}) .$$

Since  $N(B) = \{0\}$ , Corollary 5.2.4 assures that  $\mathcal{S}(\mathcal{F}) = \{z\}$ , and  $J_0 = \{1, 2\}$  is the maximal support for  $\mathcal{S}(\mathcal{F})$ . Taking  $\mathbf{a} = z^{\frac{1}{2}}$ , we have that  $\mathbf{a} \cdot \mathcal{F}$  is the unique scaled sequence of  $\mathcal{F}$  with minimal Frame Potential, but it fails to be a frame for  $\mathbb{C}^3$ , because it has just two non zero elements.  $\triangle$

**Example 5.2.8.** It can be proved that every  $G \in \mathcal{M}_3(\mathbb{C})^+$  such that  $\text{rk } G = 2$  and  $G_{ii} = 1$  for every  $i \in \mathbb{I}_3$  (considered as the Gramm matrix of a frame  $\mathcal{F}$  for  $\mathbb{C}^2$  with three unitary elements), satisfies that the minimizers  $\mathbf{a} \cdot \mathcal{F}$  of the BF-potential are frames for  $\mathbb{C}^2$ .

Indeed, given  $z \in \mathcal{S}(\mathcal{F})$ , it is easy to see that  $J = \text{supp}\{z\}$  has more than one element (otherwise  $z = e_i$  for some  $i \in \mathbb{I}_3$ ). If  $J = \mathbb{I}_3$  there is nothing to prove. Assume that  $\text{supp}\{z\} = J$  with  $|J| = 2$ . If  $\text{rk } G_J < 2$  we must have  $B_J = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ . In this case,  $I(B_J) = I_{sp}(B) = 1$ . But the

unique matrix  $B \in \mathcal{M}_3(\mathbb{C})^+$  which satisfies that  $0 \leq B$ ,  $I_{sp}(B) = 1$  and  $B_{ii} = 1$  for every  $i \in \mathbb{I}_3$  is  $B = \mathbb{1} \cdot \mathbb{1}^*$ . Indeed, if some  $B_{ij} < 1$ , then  $I_{sp}(B) \leq I_{sp} \begin{pmatrix} 1 & B_{ij} \\ B_{ij} & 1 \end{pmatrix} = \frac{1+B_{ij}}{2} < 1$ . Finally, since  $1 = \text{rk } \mathbb{1} \cdot \mathbb{1}^* = \text{rk } B = \text{rk } G \circ \overline{G} \geq \text{rk } G = 2$ , we have a contradiction.  $\triangle$

**Remark 5.2.9.** Let  $\mathcal{W} = \{W_i\}_{i \in \mathbb{I}_m}$  be a generating set of subspaces. Given a partition  $\{J_k\}_{k \in \mathbb{I}_p}$  of the set  $\mathbb{I}_m$ , we say that the sequence  $\{\mathcal{W}_k\}_{k \in \mathbb{I}_p}$  of  $\mathcal{W}$  given by  $\mathcal{W}_k = (W_i)_{i \in J_k}$  is a *partition in orthogonal components* of  $\mathcal{W}$  (briefly POC) if  $W_i \perp W_j$  for every pair  $i \in J_k, j \in \mathbb{I}_m \setminus J_k$ .

Note that by definition the trivial partition given by  $J_1 = \mathbb{I}_m$  produces a POC of  $\mathcal{W}$ . If  $\{\mathcal{W}_k\}_{k \in \mathbb{I}_p}$  is a POC of  $\mathcal{W}$ , we say that it is *maximal* if the only POC of each  $\mathcal{W}_k$  is the trivial one. It is clear that there always exists such a maximal POC for  $\mathcal{W}$ .

Let  $\{\mathcal{W}_k\}_{k \in \mathbb{I}_p}$  be a maximal POC of  $\mathcal{W}$  with  $|J_k| = m_k$  for  $1 \leq k \leq p$ . Let  $\mathbf{a}_k \in \mathbb{R}^{m_k}$  be such that  $\|\mathbf{a}_k\| = 1$  and  $I(\mathcal{W}_k) = \text{FFP}(\mathbf{a}_k \cdot \mathcal{W}_k)$  for each  $1 \leq k \leq p$ . Then, there exists  $\gamma = (\gamma_k)_{k \in \mathbb{I}_p} \in \mathbb{R}_{>0}^p$  with  $\|\gamma\| = 1$  and such that

$$I(\mathcal{W}) = \sum_{k=1}^p \text{FFP}(\gamma_k \mathbf{a}_k \cdot \mathcal{W}_k).$$

Conversely, if  $\mathbf{a} = (\mathbf{a}_1, \dots, \mathbf{a}_p)$  with  $0 \leq \mathbf{a}_k \in \mathbb{R}^{m_k}$  and  $\|\mathbf{a}\| = 1$  is such that  $I(\mathcal{W}) = \text{FFP}(\mathbf{a} \cdot \mathcal{W})$  then  $\mathbf{a}_k \neq 0$  and  $I(\mathcal{W}_k) = \text{FFP}(\|\mathbf{a}_k\|^{-1} \mathbf{a}_k \cdot \mathcal{W}_k)$ , for  $1 \leq k \leq p$ . Hence, we can restrict our study of the optimal weight of sequence of subspaces to each of the components of the maximal partition. This in turn implies that we can reduce the problem of describing the optimal weights to the case where the matrix  $B$  (which has non-negative entries and is positive semi-definite) is irreducible i.e., none of its symmetric permutations can be written as the direct sum of two matrices. This last property is relevant in the theory of matrices with non-negative entries.

## A Hadamard products and indexes.

In this section we recall some definitions and results from [12] which are closely related with the problems of Section 4. The exposition is done with some detail for several reasons: a) Most results we state are explicitly used in the previous section. b) The formulation of these results given in [12] is quite technical and intricate, so we intend here to give a clarified version. c) Although some results in the appendix are not directly applied, they are included since they give effective criteria for computing the indexes and the vectors that realize them. This is relevant since we have identified these objects as the optimal weights and the minimal potential for fusion frames.

### A.1 Basic definitions and properties

We begin with an extended version of Definition 5.1.4

**Definition A.1.1.** Let  $G \in \mathcal{M}_m(\mathbb{C})^+$ .

1. The **minimal**-Hadamard index of  $G$  is the number

$$I(G) = \max\{\lambda \geq 0 : G \circ B \geq \lambda B \quad \text{for every } B \in \mathcal{M}_m(\mathbb{C})^+\}.$$

2. Given an u.i.n  $N$  in  $\mathcal{M}_m(\mathbb{C})$ , the  $N$ -Hadamard index of  $G$  is

$$\begin{aligned} I_N(G) &= \max \{ \lambda \geq 0 : N(G \circ B) \geq \lambda N(B) \quad \text{for every } B \in \mathcal{M}_m(\mathbb{C})^+ \} \\ &= \min \{ N(G \circ B) : B \in \mathcal{M}_m(\mathbb{C})^+ \text{ and } N(B) = 1 \}. \end{aligned}$$

The index of  $G$  associated with the spectral norm  $\|\cdot\| = \|\cdot\|_{sp}$  is denoted by  $I_{sp}(G)$ , and the one associated with the Frobenius norm  $\|\cdot\|_2$  is denoted by  $I_2(G)$ .  $\triangle$

**Proposition A.1.2.** Let  $G \in \mathcal{M}_m(\mathbb{C})^+$ ,  $\mathbf{1} = (1, 1, \dots, 1) \in \mathbb{C}^m$  and  $\mathbb{E} = \mathbf{1} \cdot \mathbf{1}^T$ .

1.  $I(G) \neq 0$  if and only if  $\mathbf{1} \in R(G)$ . If there exists  $y \in \mathbb{C}^n$  such that  $Gy = \mathbf{1}$ , then

$$I(G)^{-1} = \sum_{i \in \mathbb{I}_m} y_i = \langle y, \mathbf{1} \rangle = \rho(G^\dagger \mathbb{E}) = \min \{ \langle Gz, z \rangle : \sum_{i \in \mathbb{I}_m} z_i = 1 \} . \quad (31)$$

$$\text{If } G > 0, \text{ then also } I(G) = \left( \sum_{i,j=1}^n (G^{-1})_{ij} \right)^{-1} = \frac{\det G}{\det(G + \mathbb{E}) - \det G} .$$

2.  $I(G) \leq I_N(G)$  for every u.i.n.  $N$ .
3. If  $J \subseteq \mathbb{I}_m$ , then  $I(G_J) \geq I(G)$  and  $I_N(G_J) \geq I_N(G)$ .
4. If  $D = \text{diag}(d) \in \mathcal{M}_m(\mathbb{C})^+$  is **diagonal**, then  $I_N(D) = N'(d^{-1})^{-1}$ . In particular,

$$I(D) = I_{sp}(D) = \left( \sum_{i \in \mathbb{I}_m} d_i^{-1} \right)^{-1} \quad \text{and} \quad I_2(D) = \left( \sum_{i \in \mathbb{I}_m} d_i^{-2} \right)^{-\frac{1}{2}} .$$

5. Both indexes  $I_2$  e  $I_{sp}$  are attained by matrices  $B \in \mathcal{M}_m(\mathbb{C})^+$  of rank one. This means that

$$I_2(G) = \min_{\|x\|=1} \|G \circ xx^*\|_2 \quad \text{and} \quad I_{sp}(G) = \min_{\|y\|=1} \|G \circ yy^*\| .$$

Moreover, the minima are also attained at vectors  $x \geq 0$  (or  $y \geq 0$ ).

6. Let  $B = G \circ \overline{G} \in \mathcal{M}_m(\mathbb{R})^+$ . Then  $I_2(G) = I_{sp}(B)^{\frac{1}{2}} = I_{sp}(G \circ \overline{G})^{\frac{1}{2}}$ .
7. Moreover, if  $0 \leq B \in \mathcal{M}_m(\mathbb{R})^+$  and  $A \in \mathcal{M}_n(\mathbb{R})_{sa}$  is given by  $A_{ij} = B_{ij}^{1/2}$  for  $1 \leq i, j \leq m$  then, even if  $A \notin \mathcal{M}_m(\mathbb{R})^+$ , the index  $I_2(A)$  of Definition 5.1.4 still satisfies

$$I_2(A) = \min_{\|x\|=1} \|A \circ xx^*\|_2 = \min_{\|x\|=1} \|B \circ xx^*\|_2^{\frac{1}{2}} . \quad (32)$$

8. It holds that  $I_{sp}(G) = \inf \{ I_{sp}(D) : G \leq D \text{ and } D \text{ is diagonal} \}$ . Therefore

$$I_2(G) = \inf \{ \left( \sum_{i=1}^n d_{ii}^{-2} \right)^{-\frac{1}{2}} : 0 < D \text{ is diagonal and } G \circ \overline{G} \leq D^2 \} . \quad \square$$

**Proposition A.1.3.** Let  $G \in \mathcal{M}_m(\mathbb{R})^+$  such that  $0 \leq G$ . Then  $I_{sp}(G) = I(G) \neq 0 \iff$  there exists  $u \geq 0$  such that  $Gu = \mathbf{1}$ .  $\square$

**Proposition A.1.4.** Let  $G \in \mathcal{M}_m(\mathbb{C})^+$ . Denote by  $P = G \circ \overline{G}$ . If  $x \in \mathbb{R}_*^m$ , then

$$\|G \circ xx^*\|_2^2 = \sum_{i,j} |G_{ij}|^2 |x_i|^2 |x_j|^2 = \langle P(x \circ x), x \circ x \rangle = \langle (P \circ xx^*)x, x \rangle \leq \|P \circ xx^*\| .$$

Take  $x \geq 0$  such that  $\|x\| = 1$  and  $\|G \circ xx^*\|_2 = I_2(G)$ . Then

$$(G \circ \overline{G} \circ xx^*)x = I(P_J)x , \quad \text{where} \quad J = \{i \in \mathbb{I}_m : x_i \neq 0\} .$$

In this case, they hold that

1. The vector  $u = I(P_J)^{-1}(x_J \circ x_J) \in \mathbb{C}^J$  has strict positive entries and  $P_J u = \mathbf{1}_J$ .
2.  $I_{sp}(P) = I_{sp}(P_J) = I(P_J)$ .
3.  $I_{sp}(P) = \|P \circ xx^*\|$ .  $\square$



## References

- [1] T. Ando, *Majorization, doubly stochastic matrices and comparison of eigenvalues*, Lecture Notes, Hokkaido Univ., 1982.
- [2] J.J. Benedetto and M. Fickus, *Finite normalized tight frames*, Adv. Comput. Math. **18**, 2003. 357-385.
- [3] R. Bhatia, *Matrix analysis*. Graduate Texts in Mathematics 169, Springer-Verlag, New York (1997).
- [4] B.G. Bodmann, *Optimal linear transmission by loss-insensitive packet encoding* Appl. Comput. Harmon. Anal. **22** (2007), no. 3, 274-285.
- [5] B.G. Bodmann, D.W. Kribs, V.I. Paulsen *Decoherence-Insensitive Quantum Communication by Optimal  $C^*$ -Encoding*, IEEE Transactions on Information Theory **53** (2007) 4738-4749.
- [6] P.G. Casazza, M. Fickus, J. Kovacevic, M.T. Leon, J.C. Tremain, *A physical interpretation for tight frames*, Harmonic analysis and applications, 51-76, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 2006.
- [7] P. G. Casazza and M. Fickus, *Minimizing fusion frame potential*, Acta Appl. Math., to appear.
- [8] P.G. Casazza and G. Kutyniok, *Frames of subspaces*, in “Wavelets, Frames and Operator Theory” (College Park, MD, 2003), Contemp. Math. **345**, Amer. Math. Soc., Providence, RI, 2004, 87-113.
- [9] P. G. Casazza and G. Kutyniok, *Robustness of Fusion Frames under Erasures of Subspaces and of Local Frame Vectors*, Contemp. Math., Amer. Math. Soc., Providence, RI, to appear.
- [10] P. G. Casazza, G. Kutyniok, and S. Li, *Fusion Frames and Distributed Processing*, Appl. Comput. Harmon. Anal. **25** (2008), 114-132.
- [11] P. G. Casazza, G. Kutyniok, S. Li, and C. J. Rozell, *Modeling Sensor Networks with Fusion Frames*, Wavelets XII (San Diego, CA, 2007), 67011M-1 - 67011M-11, SPIE Proc. 6701, SPIE, Bellingham, WA, 2007.
- [12] G. Corach and D. Stojanoff, *Index of Hadamard multiplication by positive matrices II*, Proceedings of the Eighth Conference of the International Linear Algebra Society (Barcelona, 1999), Linear Algebra Appl. **332/334** (2001), 503-517.
- [13] W. Fulton, *Young tableaux. With applications to representation theory and geometry* London Mathematical Society Student Texts, **35**. Cambridge University Press, Cambridge, (1997).
- [14] W. Fulton, *Eigenvalues, invariant factors, highest weights, and Schubert calculus* Bull. Amer. Math. Soc. (N.S.) **37** (2000), no. 3, 209-249.
- [15] A. Klyachko, *Stable bundles, representation theory and Hermitian operators*, Selecta Math. (N.S.) **4** (1998), no. 3, 419-445.
- [16] G. Kutyniok, A. Pezeshki, R. Calderbank and T. Liu, *Robust Dimension Reduction, Fusion Frames, and Grassmannian Packings*, Appl. Comput. Harmon. Anal. **26**, Issue 1, January 2009, 64-76 .
- [17] P. Massey, *Optimal reconstruction systems for erasures and the  $q$ -potential*, preprint.
- [18] P. Massey and M. Ruiz *Minimization of convex functionals over frame operators*, Adv. Comput. Math. (2008) doi: 10.1007/s10444-008-9092-5.
- [19] C.J. Rozell and D.H. Johnson, *Analyzing the robustness of redundant population codes in sensory and feature extraction systems*, Neurocomputing, **69** (2006) 1215-1218.
- [20] M. Ruiz and D. Stojanoff, *Some properties of frames of subspaces obtained by operator theory methods*, J. Math. Anal. Appl. **343** (2008), no. 1, 366-378.

**Pedro Massey, Mariano Ruiz and Demetrio Stojanoff**

Depto. de Matemática, FCE-UNLP, La Plata, Argentina and IAM-CONICET

e-mail: massey@mate.unlp.edu.ar , maruiz@mate.unlp.edu.ar and demetrio@mate.unlp.edu.ar