

OPERATOR INEQUALITIES RELATED TO THE CORACH–PORTA–RECHT INEQUALITY

CRISTIAN CONDE¹, MOHAMMAD SAL MOSLEHIAN², AMEUR SEDDIK³

ABSTRACT. Using elementary techniques we prove that if A, B are invertible positive operators in $\mathbb{B}(\mathcal{H})$, $t \leq 2$ and $r \in [\frac{1}{2}, \frac{3}{2}]$, then

$$(2+t) \left\| \|A^r X B^{2-r} + A^{2-r} X B^r\| \right\| \leq 2 \left\| \|A^2 X + t A X B + X B^2\| \right\|$$

for any unitarily invariant norm $\|\cdot\|$ and X in the associated ideal $\mathfrak{J}_{\|\cdot\|}$. We also characterize the class of operators satisfying $\|S X S^{-1} + S^{-1} X S + k X\| \geq (k+2) \|X\|$ under certain conditions.

1. INTRODUCTION

Let $\mathbb{B}(\mathcal{H})$, $\mathfrak{I}(\mathcal{H})$ and $\mathfrak{U}(\mathcal{H})$ be the C^* -algebra of all bounded linear operators acting on a complex Hilbert space \mathcal{H} , the set of all invertible elements in $\mathbb{B}(\mathcal{H})$ and the class of all unitary operators in $\mathbb{B}(\mathcal{H})$, respectively. The operator norm on $\mathbb{B}(\mathcal{H})$ is denoted by $\|\cdot\|$. We denote by

- $\mathcal{S}_0(\mathcal{H})$, the set of all invertible self-adjoint operators in $\mathbb{B}(\mathcal{H})$,
- $\mathcal{P}(\mathcal{H})$, the set of all positive operators in $\mathbb{B}(\mathcal{H})$,
- $\mathcal{P}_0(\mathcal{H})$, the set of all invertible positive operators in $\mathbb{B}(\mathcal{H})$,
- $\mathfrak{U}_r(\mathcal{H}) = \mathcal{S}_0(\mathcal{H}) \cap \mathfrak{U}(\mathcal{H})$, the set of all unitary reflection operators in $\mathbb{B}(\mathcal{H})$,
- $\mathfrak{N}_0(\mathcal{H})$, the set of all invertible normal operators in $\mathbb{B}(\mathcal{H})$.

For $1 \leq p < \infty$, the Schatten p -norm class consists of all compact operators A for which $\|A\|_p := (\text{tr}|A|^p)^{1/p} < \infty$, where tr is the usual trace functional. If A and B are operators in $\mathbb{B}(\mathcal{H})$ we use $A \oplus B$ to denote the 2×2 operator matrix $\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$, regarded as an operator on $\mathcal{H} \oplus \mathcal{H}$. One can show that

$$\|A \oplus B\| = \max(\|A\|, \|B\|), \quad \|A \oplus B\|_p = (\|A\|_p^p + \|B\|_p^p)^{1/p} \quad (1.1)$$

2010 *Mathematics Subject Classification*. Primary 47B47; Secondary 47A63, 47A30.

Key words and phrases. Invertible operator, unitarily invariant norm, Heinz inequality, Corach–Porta–Recht inequality, operator inequality.

One of the most essential inequalities in the operator theory is the following so-called Heinz inequality:

$$\forall P, Q \in \mathcal{P}(\mathcal{H}) \forall X \in \mathbb{B}(\mathcal{H}) \forall \alpha \in [0, 1], \quad \|PX + XQ\| \geq \|P^\alpha X Q^{1-\alpha} + P^{1-\alpha} X Q^\alpha\| \quad (1.2)$$

The proof given by Heinz [7] is based on the complex analysis and is somewhat complicated. In [9], McIntosh showed that the Heinz inequality is a consequence of the following inequality

$$\forall A, B, X \in \mathbb{B}(\mathcal{H}), \quad \|A^*AX + XBB^*\| \geq 2\|AXB\| \quad (1.3)$$

McIntosh proved that (1.3) holds and gave his ingenious proof of (1.3) \Rightarrow (1.2). In the literature, inequality (1.3) is called ‘‘Arithmetic-geometric-Mean Inequality’’.

In [4] Corach–Porta–Recht proved the following, so-called C-P-R inequality,

$$\forall S \in \mathcal{S}_0(\mathcal{H}) \forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| \geq 2\|X\| \quad (1.4)$$

The C-P-R inequality is a key factor in their study of differential geometry of self-adjoint operators. They proved this inequality by using the integral representation of a self-adjoint operator with respect to a spectral measure.

An immediate consequence of the C-P-R inequality is the following:

$$\forall S, T \in \mathcal{S}_0(\mathcal{H}) \forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXT^{-1} + S^{-1}XT\| \geq 2\|X\| \quad (1.5)$$

Using the polar decomposition of an operator, we may deduce easily from the C-P-R inequality the following operator inequality

$$\forall S \in \mathfrak{I}(\mathcal{H}) \forall X \in \mathbb{B}(\mathcal{H}), \quad \|S^*XS^{-1} + S^{-1}XS^*\| \geq 2\|X\| \quad (1.6)$$

Three years after and in [5], Fujii–Fujii–Furuta–Nakamoto proved that inequalities (1.2), (1.3), (1.4), (1.5) and two other ones hold and are mutually equivalent. By giving an easy proof of one of them, they showed a simplified proof of Heinz inequality, see also [6]. Also, it is easy to see that two inequalities (1.4) and (1.6) are equivalent.

In [10], it is shown that the operator inequality

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| \geq 2\|X\|, \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.7)$$

is in fact a characterization of $\mathbb{C}^*\mathcal{S}_0(\mathcal{H}) = \{\lambda M : \lambda \in \mathbb{C} \setminus \{0\}, M \in \mathcal{S}_0(\mathcal{H})\}$.

Recently in [11], using inequality (1.6) and the above characterization of $\mathbb{C}^*\mathcal{S}_0(\mathcal{H})$, it is proved that this class is also characterized by each of the following statements:

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| = \|S^*XS^{-1} + S^{-1}XS^*\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.8)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| \geq \|S^*XS^{-1} + S^{-1}XS^*\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.9)$$

Note that this class of operators is the class of all invertible normal operators in $\mathbb{B}(\mathcal{H})$ the spectrum of which is included in a straight line passing through the origin.

For the class of all invertible normal operators in $\mathbb{B}(\mathcal{H})$, it is proved [11, 12] that this class is characterized by each of the following properties

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1}\| + \|S^{-1}XS\| \geq 2\|X\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.10)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1}\| + \|S^{-1}XS\| = \|S^*XS^{-1}\| + \|S^{-1}XS^*\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.11)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1}\| + \|S^{-1}XS\| \geq \|S^*XS^{-1}\| + \|S^{-1}XS^*\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.12)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1}\| + \|S^{-1}XS\| \leq \|S^*XS^{-1}\| + \|S^{-1}XS^*\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.13)$$

It is natural to ask what happen if we consider in each of the above operator inequalities instead of “ \geq ”, either “ \leq ” or “ $=$ ”.

Let us consider the following associated operator inequalities

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| \leq 2\|X\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.14)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| = 2\|X\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.15)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS\| \leq \|S^*XS^{-1} + S^{-1}XS^*\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.16)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1}\| + \|S^{-1}XS\| = 2\|X\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.17)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1}\| + \|S^{-1}XS\| \leq 2\|X\| \quad (S \in \mathfrak{I}(\mathcal{H})) \quad (1.18)$$

In [11, 12, 13], it was established that each of inequalities (1.14), (1.17) and (1.18) characterize $\mathbb{R}^*\mathfrak{U}(\mathcal{H})$ and (1.15) characterizes $\mathbb{C}^*\mathfrak{U}_r(\mathcal{H})$.

We found also in [11, 12] that $\mathbb{R}^*\mathfrak{U}(\mathcal{H})$ is also characterized by each of the following two operator equalities

$$\forall X \in \mathbb{B}(\mathcal{H}), \|S^*XS^{-1} + S^{-1}XS^*\| = 2\|X\| \quad (S \in \mathfrak{J}(\mathcal{H})) \quad (1.19)$$

$$\forall X \in \mathbb{B}(\mathcal{H}), \|S^*XS^{-1}\| + \|S^{-1}XS^*\| = 2\|X\| \quad (S \in \mathfrak{J}(\mathcal{H})) \quad (1.20)$$

A unitarily invariant norm $|||\cdot|||$ is defined on a norm ideal $\mathfrak{J}_{|||\cdot|||}$ of $\mathbb{B}(\mathcal{H})$ associated with it and has the property $|||UXV||| = |||X|||$, where U and V are unitaries and $X \in \mathfrak{J}_{|||\cdot|||}$. Note that inequalities (1.2), (1.3), (1.4), (1.5) were generalized for arbitrary unitarily invariant norms. Furthermore, it is proved in [3] that the characterization of the invertible normal operators via inequalities of the uniform norm in $\mathbb{B}(\mathcal{H})$ ((1.10)-(1.14)) also holds for any unitarily invariant norm.

In [16] and in the case $\dim H < \infty$, by introducing two parameters r and t , Zhan proved that for $n \times n$ positive matrices A, B , arbitrary $n \times n$ matrix X and $(t, r) \in (-2, 2] \times [\frac{1}{2}, \frac{3}{2}]$, the following inequality holds

$$(2+t) |||A^rXB^{2-r} + A^{2-r}XB^r||| \leq 2 |||A^2X + tAXB + XB^2||| \quad (1.21)$$

for any unitarily invariant norm $|||\cdot|||$. The tool used for proving this inequality is based on the induced Schur product norm. It should be noted that the case $r = 1, t = 0$ of this result is the well-known arithmetic-geometric mean inequality due to Bhatia and Davis [1]. In this paper we want to extend it and to obtain refinements of this inequality to the case where \mathcal{H} is a Hilbert space of arbitrary dimension by using elementary techniques. We also characterize the class of operators satisfying $\|SXS^{-1} + S^{-1}XS + kX\| \geq (k+2)\|X\|$ under certain conditions.

Recently, Kittaneh proved in [8] the following refinement of the Heinz inequality.

Proposition 1.1. *Let $A, B \in \mathcal{P}(\mathcal{H})$ and $X \in \mathfrak{J}_{|||\cdot|||}$. Then*

(1) *for $\alpha \in [0, \frac{1}{2}]$ the following inequalities hold*

$$\begin{aligned} |||A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha||| &\leq |||A^{\alpha/2}XB^{1-\alpha/2} + A^{1-\alpha/2}XB^{\alpha/2}||| \\ &\leq \frac{1}{\alpha} \int_0^\alpha |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu||| d\nu \\ &\leq \frac{1}{2} |||AX + XB||| + \frac{1}{2} |||A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha||| \\ &\leq |||AX + XB||| \end{aligned} \quad (1.22)$$

(2) for $\alpha \in [\frac{1}{2}, 1]$ the following inequalities hold

$$\begin{aligned}
 |||A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha||| &\leq |||A^{\frac{1+\alpha}{2}}XB^{\frac{1-\alpha}{2}} + A^{\frac{1-\alpha}{2}}XB^{\frac{1+\alpha}{2}}||| \\
 &\leq \frac{1}{1-\alpha} \int_\alpha^1 |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu||| d\nu \\
 &\leq \frac{1}{2} |||AX + XB||| + \frac{1}{2} |||A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha||| \\
 &\leq |||AX + XB||| \tag{1.23}
 \end{aligned}$$

where

$$\begin{aligned}
 |||AX + XB||| &= \lim_{\alpha \rightarrow 0} \frac{1}{\alpha} \int_0^\alpha |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu||| d\nu \\
 &= \lim_{\alpha \rightarrow 1} \frac{1}{1-\alpha} \int_\alpha^1 |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu||| d\nu.
 \end{aligned}$$

2. MAIN RESULTS

In this section, we shall prove that inequality (1.21) of Zhan follows immediately from the generalized version of the known inequalities (1.2) and (1.4) in the more general case of arbitrary complex Hilbert space.

Theorem 2.1. *Let $A, B \in \mathcal{P}(\mathcal{H})$, where \mathcal{H} is a Hilbert space of arbitrary dimension and let $t \leq 2$, $r \in [\frac{1}{2}, \frac{3}{2}]$. Then for any unitarily invariant norm $|||\cdot|||$ and for every $X \in \mathfrak{J}_{|||\cdot|||}$, the following inequalities hold*

$$\begin{aligned}
 (1) \text{ for } r \in [\frac{1}{2}, 1] \\
 2 |||A^2X + XB^2 + tAXB||| &\geq 2 |||A^2X + XB^2 + 2AXB||| - (4 - 2t) |||AXB||| \\
 &\geq 4 |||A^{\frac{3}{2}}XB^{\frac{1}{2}} + A^{\frac{1}{2}}XB^{\frac{3}{2}}||| - (4 - 2t) |||AXB||| \\
 &\geq 2 |||A^{\frac{3}{2}}XB^{\frac{1}{2}} + A^{\frac{1}{2}}XB^{\frac{3}{2}}||| + 2 |||A^rXB^{2-r} + A^{2-r}XB^r||| - (4 - 2t) |||AXB||| \\
 &\geq \frac{4}{r - \frac{1}{2}} \int_0^{r-\frac{1}{2}} |||A^{\nu+\frac{1}{2}}XB^{\frac{3}{2}-\nu} + A^{\frac{3}{2}-\nu}XB^{\nu+\frac{1}{2}}||| d\nu - (4 - 2t) |||AXB||| \\
 &\geq 4 |||A^{\frac{2r+1}{4}}XB^{\frac{7-2r}{4}} + A^{\frac{7-2r}{4}}XB^{\frac{2r+1}{4}}||| - (4 - 2t) |||AXB||| \\
 &\geq 4 |||A^rXB^{2-r} + A^{2-r}XB^r||| - (4 - 2t) |||AXB||| \\
 &\geq (t + 2) |||A^rXB^{2-r} + A^{2-r}XB^r|||. \tag{2.1}
 \end{aligned}$$

(2) for $\alpha \in [1, \frac{3}{2}]$

$$\begin{aligned}
2 \left| \left| A^2 X + X B^2 + t A X B \right| \right| &\geq 2 \left| \left| A^2 X + X B^2 + 2 A X B \right| \right| - (4 - 2t) \left| \left| A X B \right| \right| \\
&\geq 4 \left| \left| A^{\frac{3}{2}} X B^{\frac{1}{2}} + A^{\frac{1}{2}} X B^{\frac{3}{2}} \right| \right| - (4 - 2t) \left| \left| A X B \right| \right| \\
&\geq 2 \left| \left| A^{\frac{3}{2}} X B^{\frac{1}{2}} + A^{\frac{1}{2}} X B^{\frac{3}{2}} \right| \right| + 2 \left| \left| A^r X B^{2-r} + A^{2-r} X B^r \right| \right| - (4 - 2t) \left| \left| A X B \right| \right| \\
&\geq \frac{4}{\frac{3}{2} - r} \int_{r-\frac{1}{2}}^1 \left| \left| A^{\nu+\frac{1}{2}} X B^{\frac{3}{2}-\nu} + A^{\frac{3}{2}-\nu} X B^{\nu+\frac{1}{2}} \right| \right| d\nu - (4 - 2t) \left| \left| A X B \right| \right| \\
&\geq 4 \left| \left| A^{\frac{2r+3}{4}} X B^{\frac{5-2r}{4}} + A^{\frac{5-2r}{4}} X B^{\frac{2r+3}{4}} \right| \right| - (4 - 2t) \left| \left| A X B \right| \right| \\
&\geq 4 \left| \left| A^r X B^{2-r} + A^{2-r} X B^r \right| \right| - (4 - 2t) \left| \left| A X B \right| \right| \\
&\geq (t + 2) \left| \left| A^r X B^{2-r} + A^{2-r} X B^r \right| \right|. \tag{2.2}
\end{aligned}$$

Proof. Let $X \in \mathfrak{J}_{|||,|||}$ and without loss of the generality we may assume that $A, B \in \mathcal{P}_0(\mathcal{H})$. Put $\alpha = r - \frac{1}{2}$ then $0 \leq \alpha \leq 1$.

First, we consider the case $\alpha \in [0, \frac{1}{2}]$. Using Heinz inequality and its refinements (1.22) for unitarily invariant norms and considering $A^{-\frac{1}{2}} X B^{-\frac{1}{2}} \in \mathfrak{J}_{|||,|||}$ we have

$$\begin{aligned}
&\left| \left| A^{\frac{1}{2}} X B^{-\frac{1}{2}} + A^{-\frac{1}{2}} X B^{\frac{1}{2}} \right| \right| \\
&\geq \frac{1}{2} \left(\left| \left| A^{\frac{1}{2}} X B^{-\frac{1}{2}} + A^{-\frac{1}{2}} X B^{\frac{1}{2}} \right| \right| + \left| \left| A^{\alpha-\frac{1}{2}} X B^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha} X B^{\alpha-\frac{1}{2}} \right| \right| \right) \\
&\geq \frac{1}{\alpha} \int_0^\alpha \left| \left| A^{\nu-\frac{1}{2}} X B^{\frac{1}{2}-\nu} + A^{\frac{1}{2}-\nu} X B^{\nu-\frac{1}{2}} \right| \right| d\nu \\
&\geq \left| \left| A^{\frac{\alpha-1}{2}} X B^{\frac{1-\alpha}{2}} + A^{\frac{1-\alpha}{2}} X B^{\frac{\alpha-1}{2}} \right| \right| \\
&\geq \left| \left| A^{\alpha-\frac{1}{2}} X B^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha} X B^{\alpha-\frac{1}{2}} \right| \right|. \tag{2.3}
\end{aligned}$$

Since

$$\begin{aligned}
A X B^{-1} + A^{-1} X B + 2X &= A^{\frac{1}{2}} (A^{\frac{1}{2}} X B^{-\frac{1}{2}} + A^{-\frac{1}{2}} X B^{\frac{1}{2}}) B^{-\frac{1}{2}} \\
&\quad + A^{-\frac{1}{2}} (A^{\frac{1}{2}} X B^{-\frac{1}{2}} + A^{-\frac{1}{2}} X B^{\frac{1}{2}}) B^{\frac{1}{2}},
\end{aligned}$$

utilizing the generalized version of C-P-R inequality for unitarily invariant norms, we obtain

$$\left| \left| A X B^{-1} + A^{-1} X B + 2X \right| \right| \geq 2 \left| \left| A^{\frac{1}{2}} X B^{-\frac{1}{2}} + A^{-\frac{1}{2}} X B^{\frac{1}{2}} \right| \right|. \tag{2.4}$$

It follows from (2.3) and (2.4) that

$$\begin{aligned}
 |||AXB^{-1} + A^{-1}XB + 2X||| &\geq 2 \left| \left| A^{\frac{1}{2}}XB^{-\frac{1}{2}} + A^{-\frac{1}{2}}XB^{\frac{1}{2}} \right| \right| \\
 &\geq \left| \left| A^{\frac{1}{2}}XB^{-\frac{1}{2}} + A^{-\frac{1}{2}}XB^{\frac{1}{2}} \right| \right| + \left| \left| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} \right| \right| \\
 &\geq \frac{2}{\alpha} \int_0^\alpha \left| \left| A^{\nu-\frac{1}{2}}XB^{\frac{1}{2}-\nu} + A^{\frac{1}{2}-\nu}XB^{\nu-\frac{1}{2}} \right| \right| d\nu \\
 &\geq 2 \left| \left| A^{\frac{\alpha-1}{2}}XB^{\frac{1-\alpha}{2}} + A^{\frac{1-\alpha}{2}}XB^{\frac{\alpha-1}{2}} \right| \right| \\
 &\geq 2 \left| \left| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} \right| \right|. \tag{2.5}
 \end{aligned}$$

On the other hand, due to

$$AXB^{-1} + A^{-1}XB + 2X = AXB^{-1} + A^{-1}XB + tX + (2-t)X,$$

we have

$$|||AXB^{-1} + A^{-1}XB + 2X||| \leq |||AXB^{-1} + A^{-1}XB + tX||| + (2-t) |||X|||. \tag{2.6}$$

From two last inequalities (2.5) and (2.6), we obtain

$$\begin{aligned}
 2 |||AXB^{-1} + A^{-1}XB + tX||| &\geq 2 |||AXB^{-1} + A^{-1}XB + 2X||| - (4-2t) |||X||| \\
 &\geq 4 \left| \left| A^{\frac{1}{2}}XB^{-\frac{1}{2}} + A^{-\frac{1}{2}}XB^{\frac{1}{2}} \right| \right| - (4-2t) |||X||| \\
 &\geq 2 \left| \left| A^{\frac{1}{2}}XB^{-\frac{1}{2}} + A^{-\frac{1}{2}}XB^{\frac{1}{2}} \right| \right| + 2 \left| \left| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} \right| \right| - (4-2t) |||X||| \\
 &\geq \frac{4}{\alpha} \int_0^\alpha \left| \left| A^{\nu-\frac{1}{2}}XB^{\frac{1}{2}-\nu} + A^{\frac{1}{2}-\nu}XB^{\nu-\frac{1}{2}} \right| \right| d\nu - (4-2t) |||X||| \\
 &\geq 4 \left| \left| A^{\frac{\alpha-1}{2}}XB^{\frac{1-\alpha}{2}} + A^{\frac{1-\alpha}{2}}XB^{\frac{\alpha-1}{2}} \right| \right| - (4-2t) |||X||| \\
 &\geq 4 \left| \left| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} \right| \right| - (4-2t) |||X|||. \tag{2.7}
 \end{aligned}$$

From the generalized version of C-P-R inequality for unitarily invariant norms, it is easy to see that if $s \in \mathbb{R}$

$$4 |||A^sXB^{-s} + A^{-s}XB^s||| - 4 |||X||| + 2t |||X||| \geq (t+2) |||A^sXB^{-s} + A^{-s}XB^s|||.$$

From (2.7) and the last inequality, we can deduce that for any $X \in \mathfrak{J}_{||\cdot||}$

$$\begin{aligned}
2 ||| AXB^{-1} + A^{-1}XB + tX ||| &\geq 2 ||| AXB^{-1} + A^{-1}XB + 2X ||| - (4 - 2t) ||| X ||| \\
&\geq 4 ||| A^{\frac{1}{2}}XB^{-\frac{1}{2}} + A^{-\frac{1}{2}}XB^{\frac{1}{2}} ||| - (4 - 2t) ||| X ||| \\
&\geq 2 ||| A^{\frac{1}{2}}XB^{-\frac{1}{2}} + A^{-\frac{1}{2}}XB^{\frac{1}{2}} ||| + 2 ||| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} ||| - (4 - 2t) ||| X ||| \\
&\geq \frac{4}{\alpha} \int_0^\alpha ||| A^{\nu-\frac{1}{2}}XB^{\frac{1}{2}-\nu} + A^{\frac{1}{2}-\nu}XB^{\nu-\frac{1}{2}} ||| d\nu - (4 - 2t) ||| X ||| \\
&\geq 4 ||| A^{\frac{\alpha-1}{2}}XB^{\frac{1-\alpha}{2}} + A^{\frac{1-\alpha}{2}}XB^{\frac{\alpha-1}{2}} ||| - (4 - 2t) ||| X ||| \\
&\geq 4 ||| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} ||| - (4 - 2t) ||| X ||| \\
&\geq (t + 2) ||| A^{\alpha-\frac{1}{2}}XB^{\frac{1}{2}-\alpha} + A^{\frac{1}{2}-\alpha}XB^{\alpha-\frac{1}{2}} |||. \tag{2.8}
\end{aligned}$$

whence, by replace X by AXB and α by $r - \frac{1}{2}$, we get

$$\begin{aligned}
2 ||| A^2X + XB^2 + tAXB ||| &\geq 2 ||| A^2X + XB^2 + 2AXB ||| - (4 - 2t) ||| AXB ||| \\
&\geq 4 ||| A^{\frac{3}{2}}XB^{\frac{1}{2}} + A^{\frac{1}{2}}XB^{\frac{3}{2}} ||| - (4 - 2t) ||| AXB ||| \\
&\geq 2 ||| A^{\frac{3}{2}}XB^{\frac{1}{2}} + A^{\frac{1}{2}}XB^{\frac{3}{2}} ||| + 2 ||| A^rXB^{2-r} + A^{2-r}XB^r ||| - (4 - 2t) ||| AXB ||| \\
&\geq \frac{4}{r - \frac{1}{2}} \int_0^{r-\frac{1}{2}} ||| A^{\nu+\frac{1}{2}}XB^{\frac{3}{2}-\nu} + A^{\frac{3}{2}-\nu}XB^{\nu+\frac{1}{2}} ||| d\nu - (4 - 2t) ||| AXB ||| \\
&\geq 4 ||| A^{\frac{2r+1}{4}}XB^{\frac{7-2r}{4}} + A^{\frac{7-2r}{4}}XB^{\frac{2r+1}{4}} ||| - (4 - 2t) ||| AXB ||| \\
&\geq 4 ||| A^rXB^{2-r} + A^{2-r}XB^r ||| - (4 - 2t) ||| AXB ||| \\
&\geq (t + 2) ||| A^rXB^{2-r} + A^{2-r}XB^r |||. \tag{2.9}
\end{aligned}$$

Finally, we note that the case $\alpha \in [\frac{1}{2}, 1]$ is obtained analogously and this completes the proof. \square

Note that the case $t \leq -2$ is trivial. An immediate consequence for the case $r = 1$ of this last theorem is the following (exactly the corollary 7 in [16] in finite dimensional case).

Corollary 2.2. *Let $A, B \in \mathbb{B}(\mathcal{H})$ and let $t \leq 2$. Then*

$$\forall X \in \mathfrak{J}_{||\cdot||}, ||| A^*AX + XBB^* + t|A|X|B| ||| \geq (t + 2) ||| AXB^* |||. \tag{2.10}$$

Another immediate consequence of this last corollary is (exactly the corollary 8 in [16] in finite dimensional case).

Corollary 2.3. *Let $P, Q \in \mathcal{P}_0(\mathcal{H})$ and let $t \leq 2$. Then*

$$\forall X \in \mathfrak{J}_{||\cdot||}, \quad |||PXQ^{-1} + P^{-1}XQ + tX||| \geq (t+2) |||X||| \quad (2.11)$$

Remark 2.4. The last theorem and their two consequences was proved by Zhan in [16] in the particular case of finite dimensional case. Note that Cano–Mosconi–Stojanoff [2] have proved the last corollary using the spectral measure of a normal operator to generalize [16, Corollary 8] of Zhan for arbitrary complex Hilbert space. Here, we have proved it in a general situation for an arbitrary Hilbert space using only known operator inequalities.

Remark 2.5. It follows from the above corollary that for every $k \leq 2$ and for every operator $S \in \mathbb{C}^*\mathcal{P}_0(\mathcal{H})$ the following inequality holds

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \|SXS^{-1} + S^{-1}XS + kX\| \geq (k+2) \|X\| \quad (2.12)$$

So it is interesting to characterize the class of all operators S in $\mathfrak{J}(\mathcal{H})$ satisfying this last inequality. We denote this class by $\mathfrak{D}_k(\mathcal{H})$.

Proposition 2.6. *For every real numbers k, t ,*

- (i) *if $k \geq t$, then $\mathfrak{D}_k(\mathcal{H}) \subset \mathfrak{D}_t(\mathcal{H})$,*
- (ii) *if $k \geq 0$, then*

$$\mathfrak{D}_k(\mathcal{H}) \subset \left\{ \alpha S : \alpha \in \mathbb{C}^*, S \in \mathcal{S}_0(\mathcal{H}), \left| \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right| \geq k+2, \lambda, \mu \in \sigma(S) \right\}.$$

Proof. (i) This follows by the same argument as in the proof of Theorem 2.1.

(ii) Let $S \in \mathfrak{D}_k(\mathcal{H})$. It follows immediately that the inequality $\|SXS^{-1} + S^{-1}XS\| \geq 2\|X\|$ holds for every X in $\mathbb{B}(\mathcal{H})$. Thus $S \in \mathbb{C}^*\mathcal{S}_0(\mathcal{H})$.

We may assume without loss of the generality that S is invertible and self-adjoint. Denote by $\varphi_{S,k}$ the operator on $\mathbb{B}(\mathcal{H})$ given by $\varphi_{S,k}(X) = SXS^{-1} + S^{-1}XS + kX$. So that $\sigma(\varphi_{S,k}) = \left\{ \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k : \lambda, \mu \in \sigma(S) \right\} \subset \mathbb{R}$. Hence each spectral value of $\varphi_{S,k}$ is in an approximate point value. Let $\lambda, \mu \in \sigma(S)$. Then there exists a sequence (X_n) of operators of norm one such that $\|SX_nS^{-1} + S^{-1}X_nS + kX_n\| \rightarrow \left| \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right|$. Thus $k+2 = \inf_{\|X\|=1} \|SXS^{-1} + S^{-1}XS + kX\| \leq \left| \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right|$. \square

Remark 2.7. In the case where $k \geq 0$ and $\dim H = 2$, the inclusion given in the above proposition becomes an equality.

Indeed, let S be an invertible self-adjoint operator in $\mathbb{B}(\mathcal{H})$, and let λ and μ be the eigenvalues of S such that $\left| \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right| \geq k + 2$. By a simple computation, we obtain

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad SX S^{-1} + S^{-1} X S + kX = \begin{pmatrix} k+2 & \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \\ \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k & k+2 \end{pmatrix} \circ X$$

Since the matrix $\begin{pmatrix} \frac{1}{k+2} & 1 / \left(\frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right) \\ 1 / \left(\frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right) & \frac{1}{k+2} \end{pmatrix}$ is positive definite, thus using the Schur theorem, we obtain

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \left\| \begin{pmatrix} \frac{1}{k+2} & 1 / \left(\frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right) \\ 1 / \left(\frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right) & \frac{1}{k+2} \end{pmatrix} \circ X \right\| \leq \frac{1}{k+2} \|X\|$$

Therefore

$$\forall X \in \mathbb{B}(\mathcal{H}), \quad \left\| \begin{pmatrix} k+2 & \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \\ \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k & k+2 \end{pmatrix} \circ X \right\| \geq (k+2) \|X\|$$

Conjecture 2.8. Let k be a real number such that $0 \leq k \leq 2$. Then for every natural number n and for every nonzero numbers $\lambda_1, \dots, \lambda_n$ such that $\left| \frac{\lambda_i}{\lambda_j} + \frac{\lambda_j}{\lambda_i} + k \right| \geq k + 2$, for $i, j = 1, \dots, n$ the matrix $\left(\frac{\lambda_i \lambda_j}{\lambda_i^2 + \lambda_j^2 + k \lambda_i \lambda_j} \right)_{1 \leq i, j \leq n}$ is positive.

When this conjecture is valid, the following holds;

Theorem 2.9. Assume $\dim H = n$. Then for every number k such that $0 \leq k \leq 2$ we have $\mathfrak{D}_k(\mathcal{H}) = \left\{ \alpha S : \alpha \in \mathbb{C}^*, S \in \mathcal{S}_0(\mathcal{H}), \left| \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right| \geq k + 2, \lambda, \mu \in \sigma(S) \right\}$.

Proof. Using the above Proposition, it remains to prove that

$$\left\{ \alpha S : \alpha \in \mathbb{C}^*, S \in \mathcal{S}_0(\mathcal{H}), \left| \frac{\lambda}{\mu} + \frac{\mu}{\lambda} + k \right| \geq k + 2, \lambda, \mu \in \sigma(S) \right\} \subset \mathfrak{D}_k(\mathcal{H}).$$

This follows by using the same argument used in the Remark. \square

Finally we present new variants of C-P-R inequality.

Theorem 2.10. Let $S \in \mathfrak{I}(\mathcal{H})$ and $X, Y \in \mathfrak{I}_{|||, |||}$. The following inequality holds and is equivalent to the C-P-R inequality for unitarily invariant norms:

$$(i) \quad |||(SY S^{-1} + S^{*-1} Y S^*) \oplus (S^* X S^{*-1} + S^{-1} X S)||| \geq 2 |||X \oplus Y||| ; \quad (2.13)$$

$$(ii) \quad |||(SY S^{*-1} + S^{*-1} Y S) \oplus (S^* X S^{-1} + S^{-1} X S^*)||| \geq 2 |||X \oplus Y||| . \quad (2.14)$$

Proof. (i) Clearly $\begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix}$ is a self adjoint operator in $\mathbb{B}(\mathcal{H} \oplus \mathcal{H})$ and $\begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 0 & S^{*-1} \\ S^{-1} & 0 \end{bmatrix}$. It follows from the C-P-R inequality for unitarily invariant norms that

$$\left\| \left\| \begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix} \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix} \begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix}^{-1} + \begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix}^{-1} \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix} \begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix} \right\| \right\| \geq 2 \left\| \left\| \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix} \right\| \right\|,$$

whence

$$\left\| \left\| \begin{bmatrix} SY S^{-1} + S^{*-1} Y S^* & 0 \\ 0 & S^* X S^{*-1} + S^{-1} X S \end{bmatrix} \right\| \right\| \geq 2 \left\| \left\| \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix} \right\| \right\|,$$

which is indeed (2.13). Now assume that (2.13) holds for all $X, Y \in \mathfrak{J}_{|||, |||}$ and $S \in \mathfrak{J}(\mathcal{H})$. To get (1.3) for unitarily invariant norms, let S be self-adjoint, take $Y = X$ and use the fact that two inequalities $|||A||| \leq |||B|||$ and $|||A \oplus A||| \leq |||B \oplus B|||$, by the Fan dominance principle, are equivalence for all unitarily invariant norms (see [8]).

(ii) To get inequality (2.14), use the same argument as in (i) with the matrix $\begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix}$

and note that $|||X \oplus Y||| = \left\| \left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\| \right\|$. \square

Corollary 2.11. (i) If $S \in \mathfrak{J}(\mathcal{H})$ and $X \in \mathbb{B}(\mathcal{H})$, then the following inequality holds and is equivalent to the C-P-R inequality

$$\max\{\|SXS^{-1} + S^{*-1}XS^*\|, \|S^*XS^{*-1} + S^{-1}XS\|\} \geq 2\|X\|.$$

(ii) If $S \in \mathfrak{J}(\mathcal{H})$ and X, Y are in the Schatten p -class, then the following inequality holds and is equivalent to the C-P-R inequality for the Schatten p -norm

$$\left\| \|SXS^{-1} + S^{*-1}XS^*\|_p^p + \|S^*XS^{*-1} + S^{-1}XS\|_p^p \right\| \geq 2^{p+1} \|X\|_p^p.$$

Proof. Apply (2.13) to $Y = X$ and equalities (1.1). \square

Acknowledgment. The second author was supported by a grant from Ferdowsi University of Mashhad (No. MP90113MOS).

REFERENCES

- [1] R. Bhatia and C. Davis, *More matrix forms of the arithmetic-geometric mean inequality*, SIAM J. Matrix Anal. Appl. **14** (1993), no. 1, 132–136.
- [2] C. Cano, I. Mosconi and D. Stojanoff, *Some operator inequalities for unitarily invariant norms*, Rev. Un. Mat. Argentina **46** (2005), no. 1, 53–66.
- [3] C. Conde, *Normal operators and inequalities in norm ideals*, Linear Algebra Appl. **431** (2009), no. 10, 1774–1777.
- [4] G. Corach, R. Porta, and L. Recht, *An operator inequality*, Linear Algebra Appl. **142** (1990), 153–158.
- [5] J.I. Fujii, M. Fujii, T. Furuta, R. Nakamoto, *Norm inequalities equivalent to Heinz inequality*, Proc. Amer. Math. Soc. **118** (1993), 827–830.
- [6] M. Fujii, T. Furuta, Takayuki and R. Nakamoto, *Norm inequalities in the Corach–Porta–Recht theory and operator means*, Illinois J. Math. **40** (1996), no. 4, 527–534.
- [7] E. Heinz, *Beiträge zur störungstheorie der spectralzerlegung*, Math. Ann. **123** (1951), 415–438.
- [8] F. Kittaneh, *On the convexity of the Heinz means*, Integral Equations Operator Theory **68** (2010), no. 4, 519–527.
- [9] A. McIntosh, *Heinz inequalities and perturbation of spectral families*, Macquarie Mathematical Reports, Macquarie Univ., 1979.
- [10] A. Seddik, *Some results related to Corach–Porta–Recht inequality*, Proc. Amer. Math. Soc. **129** (2001), 3009–3015.
- [11] A. Seddik, *On the injective norm and characterization of some subclasses of normal operators by inequalities or equalities*, J. Math. Anal. Appl. **351** (2009), 277–284.
- [12] A. Seddik, *Closed operator inequalities and open problems*, Math. Inequal. Appl., **14** (2011), 147–154.
- [13] A. Seddik, *Characterization of the class of unitary operators by operator inequalities*, Linear and Multilinear Algebra, to appear.
- [14] A. Seddik, *On the numerical range and norm of elementary operators*, Linear Multilinear Algebra **52** (2004), 293–302.
- [15] J.G. Stampfli, *Normality and the numerical range of an operator*, Bull. Amer. Math. Soc. **72** (1966), 1021–1022.
- [16] X. Zhan, *Inequalities for unitarily invariant norms*, SIAM J. Matrix Anal. Appl. **20** (1998), 466–470.

³ DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, TABUK UNIVERSITY, SAUDI ARABIA
E-mail address: seddikameur@hotmail.com

² DEPARTMENT OF PURE MATHEMATICS, CENTER OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, P.O. BOX 1159, MASHHAD 91775, IRAN

E-mail address: moslehan@ferdowsi.um.ac.ir and moslehan@member.ams.org

¹ INSTITUTO DE CIENCIAS, UNIVERSIDAD NACIONAL DE GRAL. SARMIENTO, J. M. GUTIERREZ 1150, (B1613GSX) LOS POLVORINES AND INSTITUTO ARGENTINO DE MATEMÁTICA “ALBERTO P. CALDERÓN”, SAAVEDRA 15 3 PISO, (C1083ACA) BUENOS AIRES, ARGENTINA

E-mail address: `cconde@ungs.edu.com`